



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/16		A1	(11) International Publication Number: WO 98/30231												
			(43) International Publication Date: 16 July 1998 (16.07.98)												
<p>(21) International Application Number: PCT/US98/00449</p> <p>(22) International Filing Date: 7 January 1998 (07.01.98)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>60/034,905</td> <td>7 January 1997 (07.01.97)</td> <td>US</td> </tr> <tr> <td>60/055,404</td> <td>8 August 1997 (08.08.97)</td> <td>US</td> </tr> <tr> <td>60/066,029</td> <td>14 November 1997 (14.11.97)</td> <td>US</td> </tr> <tr> <td>60/065,442</td> <td>14 November 1997 (14.11.97)</td> <td>US</td> </tr> </table> <p>(71) Applicant: AMYLIN PHARMACEUTICALS, INC. [US/US]; 9373 Towne Centre Drive, San Diego, CA 92121 (US).</p> <p>(72) Inventors: BEELEY, Nigel, Robert, Arnold; 227 Loma Corta Drive, Solana Beach, CA 92075 (US). PRICKETT, Kathryn, S.; 7612 Trailbrush Terrace, San Diego, CA 92126 (US). BHAVSAR, Sunil; Apartment #7, 917 Torrance Street, San Diego, CA 92103 (US).</p> <p>(74) Agents: DUFT, Bradford, J. et al.; Lyon & Lyon LLP, First Interstate World Center, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071 (US).</p>		60/034,905	7 January 1997 (07.01.97)	US	60/055,404	8 August 1997 (08.08.97)	US	60/066,029	14 November 1997 (14.11.97)	US	60/065,442	14 November 1997 (14.11.97)	US	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
60/034,905	7 January 1997 (07.01.97)	US													
60/055,404	8 August 1997 (08.08.97)	US													
60/066,029	14 November 1997 (14.11.97)	US													
60/065,442	14 November 1997 (14.11.97)	US													
(54) Title: USE OF EXENDINS AND AGONISTS THEREOF FOR THE REDUCTION OF FOOD INTAKE															
(57) Abstract															
<p>Methods for treating conditions or disorders which can be alleviated by reducing food intake are disclosed which comprise administration of an effective amount of an exendin or an exendin agonist, alone or in conjunction with other compounds or compositions that effect satiety. The methods are useful for treating conditions or disorders, including obesity, Type II diabetes, eating disorders, and insulin-resistance syndrome. The methods are also useful for lowering the plasma glucose level, lowering the plasma lipid level, reducing the cardiac risk, reducing the appetite, and reducing the weight of subjects. Pharmaceutical compositions for use in the methods of the invention are also disclosed.</p>															

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

USE OF EXENDINS AND AGONISTS THEREOF
FOR THE REDUCTION OF FOOD INTAKE

5 This application claims the benefit of U.S.
Provisional Application No. 60/034,905, filed January 7,
1997, U.S. Provisional Application No. 60/055,404, filed
August 8, 1997, U.S. Provisional Application No.
60/066,029 filed November 14, 1997, and U.S. Provisional
Application No. 60/065,442, November 14, 1997.

10

FIELD OF THE INVENTION

The present invention relates to methods for treating
conditions or disorders which can be alleviated by
reducing food intake comprising administration of an
15 effective amount of an exendin or an exendin agonist alone
or in conjunction with other compounds or compositions
that affect satiety such as a leptin or an amylin agonist.
The methods are useful for treating conditions or
disorders, in which the reduction of food intake is of
20 value, including obesity, Type II diabetes, eating
disorders, and insulin-resistance syndrome. The methods
are also useful for lowering the plasma lipid level,
reducing the cardiac risk, reducing the appetite, and
reducing the weight of subjects. Pharmaceutical
25 compositions for use in the methods of the invention are
also disclosed.

BACKGROUND

The following description summarizes information relevant to the present invention. It is not an admission that any of the information provided herein is prior art to the presently claimed invention, nor that any of the publications specifically or implicitly referenced are prior art to that invention.

Exendin

Exendins are peptides that are found in the venom of the Gila-monster, a lizard found in Arizona, and the Mexican Beaded Lizard. Exendin-3 is present in the venom of Heloderma horridum, and exendin-4 is present in the venom of Heloderma suspectum (Eng, J., et al., J. Biol. Chem., 265:20259-62, 1990; Eng, J., et al., J. Biol. Chem., 267:7402-05, 1992). The exendins have some sequence similarity to several members of the glucagon-like peptide family, with the highest homology, 53%, being to GLP-1[7-36]NH₂ (Goke, et al., J. Biol. Chem., 268:19650-55, 1993). GLP-1[7-36]NH₂, also known as proglucagon[78-107], has an insulintropic effect, stimulating insulin secretion from pancreatic β -cells; GLP also inhibits glucagon secretion from pancreatic α -cells (Orskov, et al., Diabetes, 42:658-61, 1993; D'Alessio, et al., J. Clin. Invest., 97:133-38, 1996). GLP-1 is reported to inhibit gastric emptying (Williams B, et al., J Clin Endocrinol Metab 81 (1): 327-32, 1996; Wettergren A, et al., Dig Dis Sci 38 (4): 665-73, 1993), and gastric acid

secretion. (Schjoldager BT, et al., Dig Dis Sci 34 (5): 703-8, 1989; O'Halloran DJ, et al., J Endocrinol 126 (1): 169-73, 1990; Wettergren A, et al., Dig Dis Sci 38 (4): 665-73, 1993). GLP-1[7-37], which has an additional
5 glycine residue at its carboxy terminus, also stimulates insulin secretion in humans (Orskov, et al., Diabetes, 42:658-61, 1993). A transmembrane G-protein adenylate-cyclase-coupled receptor believed to be responsible for the insulinotropic effect of GLP-1 is reported to have
10 been cloned from a β -cell line (Thorens, Proc. Natl. Acad. Sci. USA 89:8641-45 (1992)).

Exendin-4 potently binds at GLP-1 receptors on insulin-secreting β TC1 cells, at dispersed acinar cells from guinea pig pancreas, and at parietal cells from
15 stomach; the peptide is also said to stimulate somatostatin release and inhibit gastrin release in isolated stomachs (Goke, et al., J. Biol. Chem. 268:19650-55, 1993; Schepp, et al., Eur. J. Pharmacol., 69:183-91, 1994; Eissele, et al., Life Sci., 55:629-34, 1994).

20 Exendin-3 and exendin-4 were reported to stimulate cAMP production in, and amylase release from, pancreatic acinar cells (Malhotra, R., et al., Regulatory Peptides, 41:149-56, 1992; Raufman, et al., J. Biol. Chem. 267:21432-37, 1992; Singh, et al., Regul. Pept. 53:47-59, 1994). The
25 use of exendin-3 and exendin-4 as insulinotrophic agents for the treatment of diabetes mellitus and the prevention of hyperglycemia has been proposed (Eng, U.S. Patent No. 5,424,286).

C-terminally truncated exendin peptides such as exendin[9-39], a carboxyamidated molecule, and fragments 3-39 through 9-39 have been reported to be potent and selective antagonists of GLP-1 (Goke, et al., J. Biol. Chem., 268:19650-55, 1993; Raufman, J.P., et al., J. Biol. Chem. 266:2897-902, 1991; Schepp, W., et al., Eur. J. Pharm. 269:183-91, 1994; Montrose-Rafizadeh, et al., Diabetes, 45(Suppl. 2):152A, 1996). Exendin[9-39] is said to block endogenous GLP-1 in vivo, resulting in reduced insulin secretion. Wang, et al., J. Clin. Invest., 95:417-21, 1995; D'Alessio, et al., J. Clin. Invest., 97:133-38, 1996). The receptor apparently responsible for the insulinotropic effect of GLP-1 has reportedly been cloned from rat pancreatic islet cell (Thorens, B., Proc. Natl. Acad. Sci. USA 89:8641-8645, 1992). Exendins and exendin[9-39] are said to bind to the cloned GLP-1 receptor (rat pancreatic β -cell GLP-1 receptor (Fehmann HC, et al., Peptides 15 (3): 453-6, 1994) and human GLP-1 receptor (Thorens B, et al., Diabetes 42 (11): 1678-82, 1993). In cells transfected with the cloned GLP-1 receptor, exendin-4 is reportedly an agonist, i.e., it increases cAMP, while exendin[9-39] is identified as an antagonist, i.e., it blocks the stimulatory actions of exendin-4 and GLP-1. Id.

Exendin[9-39] is also reported to act as an antagonist of the full length exendins, inhibiting stimulation of pancreatic acinar cells by exendin-3 and exendin-4 (Raufman, et al., J. Biol. Chem. 266:2897-902,

1991; Raufman, et al., J. Biol. Chem., 266:21432-37, 1992). It is also reported that exendin[9-39] inhibits the stimulation of plasma insulin levels by exendin-4, and inhibits the somatostatin release-stimulating and gastrin release-inhibiting activities of exendin-4 and GLP-1 (Kolligs, F., et al., Diabetes, 44:16-19, 1995; Eissele, et al., Life Sciences, 55:629-34, 1994).

Exendins have recently been found to inhibit gastric emptying (U.S.S.N. 08/694,954, filed August 8, 1996, which enjoys common ownership with the present invention and is hereby incorporated by reference).

Exendin [9-39] has been used to investigate the physiological relevance of central GLP-1 in control of food intake (Turton, M.D. et al. Nature 379:69-72, 1996).

GLP-1 administered by intracerebroventricular injection inhibits food intake in rats. This satiety-inducing effect of GLP-1 delivered ICV is reported to be inhibited by ICV injection of exendin [9-39] (Turton, supra). However, it has been reported that GLP-1 does not inhibit food intake in mice when administered by peripheral injection (Turton, M.D., Nature 379:69-72, 1996; Bhavsar, S.P., Soc. Neurosci. Abstr. 21:460 (188.8), 1995).

Obesity and Hypernutrition

Obesity, excess adipose tissue, is becoming increasingly prevalent in developed societies. For example, approximately 30% of adults in the U.S. were estimated to be 20 percent above desirable body weight -- an accepted measure of obesity sufficient to impact a health risk (*Harrison's Principles of Internal Medicine*

12th Edition, McGraw Hill, Inc. (1991) p. 411). The pathogenesis of obesity is believed to be multifactorial but the basic problem is that in obese subjects food intake and energy expenditure do not come into balance until there is excess adipose tissue. Attempts to reduce food intake, or hypernutrition, are usually fruitless in the medium term because the weight loss induced by dieting results in both increased appetite and decreased energy expenditure (Leibel et al., (1995) *New England Journal of Medicine* 322: 621-628). The intensity of physical exercise required to expend enough energy to materially lose adipose mass is too great for most people to undertake on a sufficiently frequent basis. Thus, obesity is currently a poorly treatable, chronic, essentially intractable metabolic disorder. Not only is obesity itself believed by some to be undesirable for cosmetic reasons, but obesity also carries serious risk of co-morbidities including, Type 2 diabetes, increased cardiac risk, hypertension, atherosclerosis, degenerative arthritis, and increased incidence of complications of surgery involving general anesthesia. Obesity due to hypernutrition is also a risk factor for the group of conditions called insulin resistance syndrome, or "syndrome X." In syndrome X, it has been reported that there is a linkage between insulin resistance and hypertension. (Watson N. and Sandler M., *Curr. Med. Res. Opin.*, 12(6):374-378 (1991); Kodama J. et al., *Diabetes Care*, 13(11):1109-1111 (1990); Lithell et al., *J. Cardiovasc. Pharmacol.*, 15 Suppl. 5:S46-S52 (1990)).

In those few subjects who do succeed in losing weight, by about 10 percent of body weight, there can be striking improvements in co-morbid conditions, most especially Type 2 diabetes in which dieting and weight loss are the primary therapeutic modality, albeit relatively ineffective in many patients for the reasons stated above. Reducing food intake in obese subjects would decrease the plasma glucose level, the plasma lipid level, and the cardiac risk in these subjects.

Hypernutrition is also the result of, and the psychological cause of, many eating disorders. Reducing food intake would also be beneficial in the treatment of such disorders.

Thus, it can be appreciated that an effective means to reduce food intake is a major challenge and a superior method of treatment would be of great utility. Such a method, and compounds and compositions which are useful therefor, have been invented and are described and claimed herein.

20

SUMMARY OF THE INVENTION

The present invention concerns the surprising discovery that exendins and exendin agonists have a profound and prolonged effect on inhibiting food intake.

25

The present invention is directed to novel methods for treating conditions or disorders associated with hypernutrition, comprising the administration of an exendin, for example, exendin-3 [SEQ ID NO. 1: His Ser Asp

Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu
Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro
Ser Ser Gly Ala Pro Pro Pro Ser], or exendin-4 [SEQ ID NO.
2: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser], or other
compounds which effectively bind to the receptor at which
exendin exerts its action on reducing food intake. These
methods will be useful in the treatment of, for example,
10 obesity, diabetes, including Type II or non-insulin
dependent diabetes, eating disorders, and insulin-
resistance syndrome.

In a first aspect, the invention features a method of
treating conditions or disorders which can be alleviated
15 by reducing food intake in a subject comprising
administering to the subject a therapeutically effective
amount of an exendin or an exendin agonist. By an "exendin
agonist" is meant a compound that mimics the effects of
exendin on the reduction of food intake by binding to the
20 receptor or receptors where exendin causes this effect.
Preferred exendin agonist compounds include those described
in United States Provisional Patent Application Serial No.
60/055,404, entitled, "Novel Exendin Agonist Compounds,"
filed August 8, 1997; United States Provisional Patent
25 Application Serial No. 60/065,442, entitled, "Novel Exendin
Agonist Compounds," filed November 14, 1997; and United
States Provisional Patent Application Serial No.
60/066,029, entitled, "Novel Exendin Agonist Compounds,"
filed November 14, 1997; all of which enjoy common

ownership with the present application and all of which are incorporated by this reference into the present application as though fully set forth herein. By "condition or disorder which can be alleviated by reducing food intake" is meant any condition or disorder in a subject that is either caused by, complicated by, or aggravated by a relatively high food intake, or that can be alleviated by reducing food intake. Such conditions or disorders include, but are not limited to, obesity, diabetes, including Type II diabetes, eating disorders, and insulin-resistance syndrome.

Thus, in a first embodiment, the present invention provides a method for treating conditions or disorders which can be alleviated by reducing food intake in a subject comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist. Preferred exendin agonist compounds include those described in U.S. Provisional Patent Application Serial Nos. 60/055,404; 60/065,442; and 60/066,029, which have been incorporated by reference in the present application. Preferably, the subject is a vertebrate, more preferably a mammal, and most preferably a human. In preferred aspects, the exendin or exendin agonist is administered parenterally, more preferably by injection. In a most preferred aspect, the injection is a peripheral injection. Preferably, about 10 μ g-30 μ g to about 5 mg of the exendin or exendin agonist is administered per day. More preferably, about 10-30 μ g to about 2mg, or about 10-30 μ g to about 1mg of the exendin or

exendin agonist is administered per day. Most preferably, about 30 μ g to about 500 μ g of the exendin or exendin agonist is administered per day.

5 In various preferred embodiments of the invention, the condition or disorder is obesity, diabetes, preferably Type II diabetes, an eating disorder, or insulin-resistance syndrome.

10 In other preferred aspects of the invention, a method is provided for reducing the appetite of a subject comprising administering to said subject an appetite-lowering amount of an exendin or an exendin agonist.

15 In yet other preferred aspects, a method is provided for lowering plasma lipids comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist.

The methods of the present invention may also be used to reduce the cardiac risk of a subject comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist. In one preferred aspect, the exendin or exendin agonist used in the methods of the present invention is exendin-3. In another preferred aspect, said exendin is exendin-4. Other preferred exendin agonists include exendin-4 (1-30) [SEQ ID NO 6: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly], exendin-4 (1-30) amide [SEQ ID NO 7: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly-NH₂], exendin-4 (1-28) amide [SEQ ID NO 40: His Gly Glu Gly Thr

20

25

Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg
Leu Phe Ile Glu Trp Leu Lys Asn-NH₂], ¹⁴Leu, ²⁵Phe exendin-4
amide [SEQ ID NO 9: His Gly Glu Gly Thr Phe Thr Ser Asp Leu
Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe
5 Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-
NH₂], ¹⁴Leu, ²⁵Phe exendin-4 (1-28) amide [SEQ ID NO 41: His
Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu
Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂], and
¹⁴Leu, ²²Ala, ²⁵Phe exendin-4 (1-28) amide [SEQ ID NO 8: His Gly
10 Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu
Ala Val Arg Leu Ala Ile Glu Phe Leu Lys Asn-NH₂].

In the methods of the present invention, the exendins
and exendin agonists may be administered separately or
together with one or more other compounds and compositions
15 that exhibit a long term or short-term satiety action,
including, but not limited to other compounds and
compositions that comprise an amylin agonist,
cholecystokinin (CCK), or a leptin (ob protein). Suitable
amylin agonists include, for example, [^{25,28,29}Pro-]-human
20 amylin (also known as "pramlintide," and previously
referred to as "AC-137") as described in "Amylin Agonist
Peptides and Uses Therefor," U.S. Patent No. 5,686,511,
issued November 11, 1997, and salmon calcitonin. The CCK
used is preferably CCK octopeptide (CCK-8). Leptin is
25 discussed in, for example, Pelleymounter, M.A., et al.
Science 269:540-43 (1995); Halaas, J.L., et al. Science
269:543-46 (1995); and Campfield, L.A., et al. Eur. J.
Pharmac. 262:133-41 (1994).

In other embodiments of the invention is provided a pharmaceutical composition for use in the treatment of conditions or disorders which can be alleviated by reducing food intake comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier. Preferably, the pharmaceutical composition comprises a therapeutically effective amount for a human subject.

The pharmaceutical composition may preferably be used for reducing the appetite of a subject, reducing the weight of a subject, lowering the plasma lipid level of a subject, or reducing the cardiac risk of a subject. Those of skill in the art will recognize that the pharmaceutical composition will preferably comprise a therapeutically effective amount of an exendin or exendin agonist to accomplish the desired effect in the subject.

The pharmaceutical compositions may further comprise one or more other compounds and compositions that exhibit a long-term or short-term satiety action, including, but not limited to other compounds and compositions that comprise an amylin agonist, CCK, preferably CCK-8, or leptin. Suitable amylin agonists include, for example, [^{25,28,29}Pro]-human amylin and salmon calcitonin.

In one preferred aspect, the pharmaceutical composition comprises exendin-3. In another preferred aspect, the pharmaceutical composition comprises exendin-4.

In other preferred aspects, the pharmaceutical compositions comprises a peptide selected from: exendin-4 (1-30), exendin-4 (1-30) amide, exendin-4 (1-28) amide,

¹⁴Leu, ²⁵Phe exendin-4 amide, ¹⁴Leu, ²⁵Phe exendin-4 (1-28) amide, and ¹⁴Leu, ²²Ala, ²⁵Phe exendin-4 (1-28) amide.

5

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of exendin-4 and GLP-1.

10

Figure 2 is a graphical depiction of the change of food intake in obese mice after intraperitoneal injection of exendin-4.

Figure 3 is a graphical depiction of the change of food intake in rats after intracerebroventricular injection of exendin-4

15

Figure 4 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of exendin-4 (1-30) ("Compound 1").

20

Figure 5 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of exendin-4 (1-30) amide ("Compound 2").

Figure 6 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of exendin-4 (1-28) amide ("Compound 3").

25

Figure 7 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of ¹⁴Leu, ²⁵Phe exendin-4 amide ("Compound 4").

Figure 8 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of ¹⁴Leu, ²⁵Phe exendin-4 (1-28) amide ("Compound 5").

Figure 9 is a graphical depiction of the change of food intake in normal mice after intraperitoneal injection of ¹⁴Leu, ²²Ala, ²⁵Phe exendin-4 (1-28) amide ("Compound 6").

Figure 10 depicts the amino acid sequences for certain exendin agonist compounds useful in the present invention [SEQ ID NOS 9-39].

DETAILED DESCRIPTION OF THE INVENTION

Exendins and exendin agonists are useful as described herein in view of their pharmacological properties. Activity as exendin agonists can be indicated by activity in the assays described below. Effects of exendins or exendin agonists on reducing food intake can be identified, evaluated, or screened for, using the methods described in the Examples below, or other methods known in the art for determining effects on food intake or appetite.

Exendin Agonist Compounds

Exendin agonist compounds are those described in U.S. Provisional Application No. 60/055,404, including compounds of the formula (I) [SEQ ID NO. 3]:

1	5	10
Xaa ₁	Xaa ₂	Xaa ₃
Gly	Thr	Xaa ₄
Xaa ₅	Xaa ₆	Xaa ₇
Xaa ₈	Xaa ₉	Xaa ₁₀
Ser	Lys	Gln
Xaa ₁₁	Glu	Glu
Glu	Ala	Val
Arg	Leu	Xaa ₁₂
Xaa ₁₃	Leu	Lys
Asn	Gly	Gly
Xaa ₁₄	Xaa ₁₅	Xaa ₁₆
Xaa ₁₇	Xaa ₁₈	Xaa ₁₉
Xaa ₂₀	Xaa ₂₁	Xaa ₂₂
Xaa ₂₃	Xaa ₂₄	Xaa ₂₅
Xaa ₂₆	Xaa ₂₇	Xaa ₂₈
Xaa ₂₉	Xaa ₃₀	Xaa ₃₁
Xaa ₃₂	Xaa ₃₃	Xaa ₃₄
Xaa ₃₅	Xaa ₃₆	Xaa ₃₇
Xaa ₃₈	Xaa ₃₉	Xaa ₄₀
Xaa ₄₁	Xaa ₄₂	Xaa ₄₃
Xaa ₄₄	Xaa ₄₅	Xaa ₄₆
Xaa ₄₇	Xaa ₄₈	Xaa ₄₉
Xaa ₅₀	Xaa ₅₁	Xaa ₅₂
Xaa ₅₃	Xaa ₅₄	Xaa ₅₅
Xaa ₅₆	Xaa ₅₇	Xaa ₅₈
Xaa ₅₉	Xaa ₆₀	Xaa ₆₁
Xaa ₆₂	Xaa ₆₃	Xaa ₆₄
Xaa ₆₅	Xaa ₆₆	Xaa ₆₇
Xaa ₆₈	Xaa ₆₉	Xaa ₇₀
Xaa ₇₁	Xaa ₇₂	Xaa ₇₃
Xaa ₇₄	Xaa ₇₅	Xaa ₇₆
Xaa ₇₇	Xaa ₇₈	Xaa ₇₉
Xaa ₈₀	Xaa ₈₁	Xaa ₈₂
Xaa ₈₃	Xaa ₈₄	Xaa ₈₅
Xaa ₈₆	Xaa ₈₇	Xaa ₈₈
Xaa ₈₉	Xaa ₉₀	Xaa ₉₁
Xaa ₉₂	Xaa ₉₃	Xaa ₉₄
Xaa ₉₅	Xaa ₉₆	Xaa ₉₇
Xaa ₉₈	Xaa ₉₉	Xaa ₁₀₀
Xaa ₁₀₁	Xaa ₁₀₂	Xaa ₁₀₃
Xaa ₁₀₄	Xaa ₁₀₅	Xaa ₁₀₆
Xaa ₁₀₇	Xaa ₁₀₈	Xaa ₁₀₉
Xaa ₁₁₀	Xaa ₁₁₁	Xaa ₁₁₂
Xaa ₁₁₃	Xaa ₁₁₄	Xaa ₁₁₅
Xaa ₁₁₆	Xaa ₁₁₇	Xaa ₁₁₈
Xaa ₁₁₉	Xaa ₁₂₀	Xaa ₁₂₁
Xaa ₁₂₂	Xaa ₁₂₃	Xaa ₁₂₄
Xaa ₁₂₅	Xaa ₁₂₆	Xaa ₁₂₇
Xaa ₁₂₈	Xaa ₁₂₉	Xaa ₁₃₀
Xaa ₁₃₁	Xaa ₁₃₂	Xaa ₁₃₃
Xaa ₁₃₄	Xaa ₁₃₅	Xaa ₁₃₆
Xaa ₁₃₇	Xaa ₁₃₈	Xaa ₁₃₉
Xaa ₁₄₀	Xaa ₁₄₁	Xaa ₁₄₂
Xaa ₁₄₃	Xaa ₁₄₄	Xaa ₁₄₅
Xaa ₁₄₆	Xaa ₁₄₇	Xaa ₁₄₈
Xaa ₁₄₉	Xaa ₁₅₀	Xaa ₁₅₁
Xaa ₁₅₂	Xaa ₁₅₃	Xaa ₁₅₄
Xaa ₁₅₅	Xaa ₁₅₆	Xaa ₁₅₇
Xaa ₁₅₈	Xaa ₁₅₉	Xaa ₁₆₀
Xaa ₁₆₁	Xaa ₁₆₂	Xaa ₁₆₃
Xaa ₁₆₄	Xaa ₁₆₅	Xaa ₁₆₆
Xaa ₁₆₇	Xaa ₁₆₈	Xaa ₁₆₉
Xaa ₁₇₀	Xaa ₁₇₁	Xaa ₁₇₂
Xaa ₁₇₃	Xaa ₁₇₄	Xaa ₁₇₅
Xaa ₁₇₆	Xaa ₁₇₇	Xaa ₁₇₈
Xaa ₁₇₉	Xaa ₁₈₀	Xaa ₁₈₁
Xaa ₁₈₂	Xaa ₁₈₃	Xaa ₁₈₄
Xaa ₁₈₅	Xaa ₁₈₆	Xaa ₁₈₇
Xaa ₁₈₈	Xaa ₁₈₉	Xaa ₁₉₀
Xaa ₁₉₁	Xaa ₁₉₂	Xaa ₁₉₃
Xaa ₁₉₄	Xaa ₁₉₅	Xaa ₁₉₆
Xaa ₁₉₇	Xaa ₁₉₈	Xaa ₁₉₉
Xaa ₂₀₀	Xaa ₂₀₁	Xaa ₂₀₂
Xaa ₂₀₃	Xaa ₂₀₄	Xaa ₂₀₅
Xaa ₂₀₆	Xaa ₂₀₇	Xaa ₂₀₈
Xaa ₂₀₉	Xaa ₂₁₀	Xaa ₂₁₁
Xaa ₂₁₂	Xaa ₂₁₃	Xaa ₂₁₄
Xaa ₂₁₅	Xaa ₂₁₆	Xaa ₂₁₇
Xaa ₂₁₈	Xaa ₂₁₉	Xaa ₂₂₀
Xaa ₂₂₁	Xaa ₂₂₂	Xaa ₂₂₃
Xaa ₂₂₄	Xaa ₂₂₅	Xaa ₂₂₆
Xaa ₂₂₇	Xaa ₂₂₈	Xaa ₂₂₉
Xaa ₂₃₀	Xaa ₂₃₁	Xaa ₂₃₂
Xaa ₂₃₃	Xaa ₂₃₄	Xaa ₂₃₅
Xaa ₂₃₆	Xaa ₂₃₇	Xaa ₂₃₈
Xaa ₂₃₉	Xaa ₂₄₀	Xaa ₂₄₁
Xaa ₂₄₂	Xaa ₂₄₃	Xaa ₂₄₄
Xaa ₂₄₅	Xaa ₂₄₆	Xaa ₂₄₇
Xaa ₂₄₈	Xaa ₂₄₉	Xaa ₂₅₀
Xaa ₂₅₁	Xaa ₂₅₂	Xaa ₂₅₃
Xaa ₂₅₄	Xaa ₂₅₅	Xaa ₂₅₆
Xaa ₂₅₇	Xaa ₂₅₈	Xaa ₂₅₉
Xaa ₂₆₀	Xaa ₂₆₁	Xaa ₂₆₂
Xaa ₂₆₃	Xaa ₂₆₄	Xaa ₂₆₅
Xaa ₂₆₆	Xaa ₂₆₇	Xaa ₂₆₈
Xaa ₂₆₉	Xaa ₂₇₀	Xaa ₂₇₁
Xaa ₂₇₂	Xaa ₂₇₃	Xaa ₂₇₄
Xaa ₂₇₅	Xaa ₂₇₆	Xaa ₂₇₇
Xaa ₂₇₈	Xaa ₂₇₉	Xaa ₂₈₀
Xaa ₂₈₁	Xaa ₂₈₂	Xaa ₂₈₃
Xaa ₂₈₄	Xaa ₂₈₅	Xaa ₂₈₆
Xaa ₂₈₇	Xaa ₂₈₈	Xaa ₂₈₉
Xaa ₂₉₀	Xaa ₂₉₁	Xaa ₂₉₂
Xaa ₂₉₃	Xaa ₂₉₄	Xaa ₂₉₅
Xaa ₂₉₆	Xaa ₂₉₇	Xaa ₂₉₈
Xaa ₂₉₉	Xaa ₃₀₀	Xaa ₃₀₁
Xaa ₃₀₂	Xaa ₃₀₃	Xaa ₃₀₄
Xaa ₃₀₅	Xaa ₃₀₆	Xaa ₃₀₇
Xaa ₃₀₈	Xaa ₃₀₉	Xaa ₃₁₀
Xaa ₃₁₁	Xaa ₃₁₂	Xaa ₃₁₃
Xaa ₃₁₄	Xaa ₃₁₅	Xaa ₃₁₆
Xaa ₃₁₇	Xaa ₃₁₈	Xaa ₃₁₉
Xaa ₃₂₀	Xaa ₃₂₁	Xaa ₃₂₂
Xaa ₃₂₃	Xaa ₃₂₄	Xaa ₃₂₅
Xaa ₃₂₆	Xaa ₃₂₇	Xaa ₃₂₈
Xaa ₃₂₉	Xaa ₃₃₀	Xaa ₃₃₁
Xaa ₃₃₂	Xaa ₃₃₃	Xaa ₃₃₄
Xaa ₃₃₅	Xaa ₃₃₆	Xaa ₃₃₇
Xaa ₃₃₈	Xaa ₃₃₉	Xaa ₃₄₀
Xaa ₃₄₁	Xaa ₃₄₂	Xaa ₃₄₃
Xaa ₃₄₄	Xaa ₃₄₅	Xaa ₃₄₆
Xaa ₃₄₇	Xaa ₃₄₈	Xaa ₃₄₉
Xaa ₃₅₀	Xaa ₃₅₁	Xaa ₃₅₂
Xaa ₃₅₃	Xaa ₃₅₄	Xaa ₃₅₅
Xaa ₃₅₆	Xaa ₃₅₇	Xaa ₃₅₈
Xaa ₃₅₉	Xaa ₃₆₀	Xaa ₃₆₁
Xaa ₃₆₂	Xaa ₃₆₃	Xaa ₃₆₄
Xaa ₃₆₅	Xaa ₃₆₆	Xaa ₃₆₇
Xaa ₃₆₈	Xaa ₃₆₉	Xaa ₃₇₀
Xaa ₃₇₁	Xaa ₃₇₂	Xaa ₃₇₃
Xaa ₃₇₄	Xaa ₃₇₅	Xaa ₃₇₆
Xaa ₃₇₇	Xaa ₃₇₈	Xaa ₃₇₉
Xaa ₃₈₀	Xaa ₃₈₁	Xaa ₃₈₂
Xaa ₃₈₃	Xaa ₃₈₄	Xaa ₃₈₅
Xaa ₃₈₆	Xaa ₃₈₇	Xaa ₃₈₈
Xaa ₃₈₉	Xaa ₃₉₀	Xaa ₃₉₁
Xaa ₃₉₂	Xaa ₃₉₃	Xaa ₃₉₄
Xaa ₃₉₅	Xaa ₃₉₆	Xaa ₃₉₇
Xaa ₃₉₈	Xaa ₃₉₉	Xaa ₄₀₀
Xaa ₄₀₁	Xaa ₄₀₂	Xaa ₄₀₃
Xaa ₄₀₄	Xaa ₄₀₅	Xaa ₄₀₆
Xaa ₄₀₇	Xaa ₄₀₈	Xaa ₄₀₉
Xaa ₄₁₀	Xaa ₄₁₁	Xaa ₄₁₂
Xaa ₄₁₃	Xaa ₄₁₄	Xaa ₄₁₅
Xaa ₄₁₆	Xaa ₄₁₇	Xaa ₄₁₈
Xaa ₄₁₉	Xaa ₄₂₀	Xaa ₄₂₁
Xaa ₄₂₂	Xaa ₄₂₃	Xaa ₄₂₄
Xaa ₄₂₅	Xaa ₄₂₆	Xaa ₄₂₇
Xaa ₄₂₈	Xaa ₄₂₉	Xaa ₄₃₀
Xaa ₄₃₁	Xaa ₄₃₂	Xaa ₄₃₃
Xaa ₄₃₄	Xaa ₄₃₅	Xaa ₄₃₆
Xaa ₄₃₇	Xaa ₄₃₈	Xaa ₄₃₉
Xaa ₄₄₀	Xaa ₄₄₁	Xaa ₄₄₂
Xaa ₄₄₃	Xaa ₄₄₄	Xaa ₄₄₅
Xaa ₄₄₆	Xaa ₄₄₇	Xaa ₄₄₈
Xaa ₄₄₉	Xaa ₄₅₀	Xaa ₄₅₁
Xaa ₄₅₂	Xaa ₄₅₃	Xaa ₄₅₄
Xaa ₄₅₅	Xaa ₄₅₆	Xaa ₄₅₇
Xaa ₄₅₈	Xaa ₄₅₉	Xaa ₄₆₀
Xaa ₄₆₁	Xaa ₄₆₂	Xaa ₄₆₃
Xaa ₄₆₄	Xaa ₄₆₅	Xaa ₄₆₆
Xaa ₄₆₇	Xaa ₄₆₈	Xaa ₄₆₉
Xaa ₄₇₀	Xaa ₄₇₁	Xaa ₄₇₂
Xaa ₄₇₃	Xaa ₄₇₄	Xaa ₄₇₅
Xaa ₄₇₆	Xaa ₄₇₇	Xaa ₄₇₈
Xaa ₄₇₉	Xaa ₄₈₀	Xaa ₄₈₁
Xaa ₄₈₂	Xaa ₄₈₃	Xaa ₄₈₄
Xaa ₄₈₅	Xaa ₄₈₆	Xaa ₄₈₇
Xaa ₄₈₈	Xaa ₄₈₉	Xaa ₄₉₀
Xaa ₄₉₁	Xaa ₄₉₂	Xaa ₄₉₃
Xaa ₄₉₄	Xaa ₄₉₅	Xaa ₄₉₆
Xaa ₄₉₇	Xaa ₄₉₈	Xaa ₄₉₉
Xaa ₅₀₀	Xaa ₅₀₁	Xaa ₅₀₂
Xaa ₅₀₃	Xaa ₅₀₄	Xaa ₅₀₅
Xaa ₅₀₆	Xaa ₅₀₇	Xaa ₅₀₈
Xaa ₅₀₉	Xaa ₅₁₀	Xaa ₅₁₁
Xaa ₅₁₂	Xaa ₅₁₃	Xaa ₅₁₄
Xaa ₅₁₅	Xaa ₅₁₆	Xaa ₅₁₇
Xaa ₅₁₈	Xaa ₅₁₉	Xaa ₅₂₀
Xaa ₅₂₁	Xaa ₅₂₂	Xaa ₅₂₃
Xaa ₅₂₄	Xaa ₅₂₅	Xaa ₅₂₆
Xaa ₅₂₇	Xaa ₅₂₈	Xaa ₅₂₉
Xaa ₅₃₀	Xaa ₅₃₁	Xaa ₅₃₂
Xaa ₅₃₃	Xaa ₅₃₄	Xaa ₅₃₅
Xaa ₅₃₆	Xaa ₅₃₇	Xaa ₅₃₈
Xaa ₅₃₉	Xaa ₅₄₀	Xaa ₅₄₁
Xaa ₅₄₂	Xaa ₅₄₃	Xaa ₅₄₄
Xaa ₅₄₅	Xaa ₅₄₆	Xaa ₅₄₇
Xaa ₅₄₈	Xaa ₅₄₉	Xaa ₅₅₀
Xaa ₅₅₁	Xaa ₅₅₂	Xaa ₅₅₃
Xaa ₅₅₄	Xaa ₅₅₅	Xaa ₅₅₆
Xaa ₅₅₇	Xaa ₅₅₈	Xaa ₅₅₉
Xaa ₅₆₀	Xaa ₅₆₁	Xaa ₅₆₂
Xaa ₅₆₃	Xaa ₅₆₄	Xaa ₅₆₅
Xaa ₅₆₆	Xaa ₅₆₇	Xaa ₅₆₈
Xaa ₅₆₉	Xaa ₅₇₀	Xaa ₅₇₁
Xaa ₅₇₂	Xaa ₅₇₃	Xaa ₅₇₄
Xaa ₅₇₅	Xaa ₅₇₆	Xaa ₅₇₇
Xaa ₅₇₈	Xaa ₅₇₉	Xaa ₅₈₀
Xaa ₅₈₁	Xaa ₅₈₂	Xaa ₅₈₃
Xaa ₅₈₄	Xaa ₅₈₅	Xaa ₅₈₆
Xaa ₅₈₇	Xaa ₅₈₈	Xaa ₅₈₉
Xaa ₅₉₀	Xaa ₅₉₁	Xaa ₅₉₂
Xaa ₅₉₃	Xaa ₅₉₄	Xaa ₅₉₅
Xaa ₅₉₆	Xaa ₅₉₇	Xaa ₅₉₈
Xaa ₅₉₉	Xaa ₆₀₀	Xaa ₆₀₁
Xaa ₆₀₂	Xaa ₆₀₃	Xaa ₆₀₄
Xaa ₆₀₅	Xaa ₆₀₆	Xaa ₆₀₇
Xaa ₆₀₈	Xaa ₆₀₉	Xaa ₆₁₀
Xaa ₆₁₁	Xaa ₆₁₂	Xaa ₆₁₃
Xaa ₆₁₄	Xaa ₆₁₅	Xaa ₆₁₆
Xaa ₆₁₇	Xaa ₆₁₈	Xaa ₆₁₉
Xaa ₆₂₀	Xaa ₆₂₁	Xaa ₆₂₂
Xaa ₆₂₃	Xaa ₆₂₄	Xaa ₆₂₅
Xaa ₆₂₆	Xaa ₆₂₇	Xaa ₆₂₈
Xaa ₆₂₉	Xaa ₆₃₀	Xaa ₆₃₁
Xaa ₆₃₂	Xaa ₆₃₃	Xaa ₆₃₄
Xaa ₆₃₅	Xaa ₆₃₆	Xaa ₆₃₇
Xaa ₆₃₈	Xaa ₆₃₉	Xaa ₆₄₀
Xaa ₆₄₁	Xaa ₆₄₂	Xaa ₆₄₃
Xaa ₆₄₄	Xaa ₆₄₅	Xaa ₆₄₆
Xaa ₆₄₇	Xaa ₆₄₈	Xaa ₆₄₉
Xaa ₆₅₀	Xaa ₆₅₁	Xaa ₆₅₂
Xaa ₆₅₃	Xaa ₆₅₄	Xaa ₆₅₅
Xaa ₆₅₆	Xaa ₆₅₇	Xaa ₆₅₈
Xaa ₆₅₉	Xaa ₆₆₀	Xaa ₆₆₁
Xaa ₆₆₂	Xaa ₆₆₃	Xaa ₆₆₄
Xaa ₆₆₅	Xaa ₆₆₆	Xaa ₆₆₇
Xaa ₆₆₈	Xaa ₆₆₉	Xaa ₆₇₀
Xaa ₆₇₁	Xaa ₆₇₂	Xaa ₆₇₃
Xaa ₆₇₄	Xaa ₆₇	

wherein Xaa₁ is His, Arg or Tyr; Xaa₂ is Ser, Gly, Ala or Thr; Xaa₃ is Asp or Glu; Xaa₄ is Phe, Tyr or naphthylalanine; Xaa₅ is Thr or Ser; Xaa₆ is Ser or Thr; Xaa₇ is Asp or Glu; Xaa₈ is Leu, Ile, Val, pentylglycine or Met; Xaa₉ is Leu, Ile, pentylglycine, Val or Met; Xaa₁₀ is Phe, Tyr or naphthylalanine; Xaa₁₁ is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met; Xaa₁₂ is Glu or Asp; Xaa₁₃ is Trp, Phe, Tyr, or naphthylalanine; Xaa₁₄, Xaa₁₅, Xaa₁₆ and Xaa₁₇ are independently Pro, homoproline, 3Hyp, 4Hyp, thioproline, N-alkylglycine, N-alkylpentylglycine or N-alkylalanine; Xaa₁₈ is Ser, Thr or Tyr; and Z is -OH or -NH₂; with the proviso that the compound is not exendin-3 or exindin-4.

Preferred N-alkyl groups for N-alkylglycine, N-alkylpentylglycine and N-alkylalanine include lower alkyl groups preferably of 1 to about 6 carbon atoms, more preferably of 1 to 4 carbon atoms. Suitable compounds include those listed in Figure 10 having amino acid sequences of SEQ. ID. NOS. 9 to 39.

Preferred exendin agonist compounds include those wherein Xaa₁ is His or Tyr. More preferably Xaa₁ is His.

Preferred are those compounds wherein Xaa₂ is Gly.

Preferred are those compounds wherein Xaa₃ is Leu, pentylglycine or Met.

Preferred compounds include those wherein Xaa₄ is Trp or Phe.

Also preferred are compounds where Xaa₄ is Phe or naphthylalanine; Xaa₁₁ is Ile or Val and Xaa₁₄, Xaa₁₅, Xaa₁₆

and Xaa₁₇, are independently selected from Pro, homoproline, thioproline or N-alkylalanine. Preferably N-alkylalanine has a N-alkyl group of 1 to about 6 carbon atoms.

According to an especially preferred aspect, Xaa₁₅,
5 Xaa₁₆ and Xaa₁₇, are the same amino acid residue.

Preferred are compounds wherein Xaa₁₈ is Ser or Tyr, more preferably Ser.

Preferably Z is -NH₂.

According to one aspect, preferred are compounds of
10 formula (I) wherein Xaa₁ is His or Tyr, more preferably His; Xaa₂ is Gly; Xaa₄ is Phe or naphthylalanine; Xaa₅ is Leu, pentylglycine or Met; Xaa₁₀ is Phe or naphthylalanine; Xaa₁₁ is Ile or Val; Xaa₁₄, Xaa₁₅, Xaa₁₆ and Xaa₁₇, are
15 independently selected from Pro, homoproline, thioproline or N-alkylalanine; and Xaa₁₈ is Ser or Tyr, more preferably Ser. More preferably Z is -NH₂.

According to an especially preferred aspect,
especially preferred compounds include those of formula
(I) wherein: Xaa₁ is His or Arg; Xaa₂ is Gly; Xaa₃ is Asp
20 or Glu; Xaa₄ is Phe or naphthylalanine; Xaa₅ is Thr or Ser;

Xaa₆ is Ser or Thr; Xaa₇ is Asp or Glu; Xaa₈ is Leu or pentylglycine; Xaa₉ is Leu or pentylglycine; Xaa₁₀ is Phe or naphthylalanine; Xaa₁₁ is Ile, Val or t-butyltylglycine; Xaa₁₂ is Glu or Asp; Xaa₁₃ is Trp or Phe; Xaa₁₄, Xaa₁₅, Xaa₁₆,
25 and Xaa₁₇, are independently Pro, homoproline, thioproline, or N-methylalanine; Xaa₁₈ is Ser or Tyr; and Z is -OH or -NH₂; with the proviso that the compound does not have the formula of either SEQ. ID. NOS. 1 or 2. More preferably Z

is -NH₂. Especially preferred compounds include those having the amino acid sequence of SEQ. ID. NOS. 9, 10, 21, 22, 23, 26, 28, 34, 35 and 39.

5 According to an especially preferred aspect, provided are compounds where Xaa₁ is Leu, Ile, Val or pentylglycine, more preferably Leu or pentylglycine, and Xaa₁₁ is Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds will exhibit advantageous
10 duration of action and be less subject to oxidative degradation, both in vitro and in vivo, as well as during synthesis of the compound.

Exendin agonist compounds also include those described in U.S. Provisional Application No. 60/065,442,
15 including compounds of the formula (II) [SEQ ID NO. 4]:

Xaa₁ Xaa₂ Xaa₃ Gly Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁; wherein

20

Xaa₁ is His, Arg or Tyr;

Xaa₂ is Ser, Gly, Ala or Thr;

Xaa₃ is Asp or Glu;

Xaa₅ is Ala or Thr;

25 Xaa₆ is Ala, Phe, Tyr or naphthylalanine;

Xaa₇ is Thr or Ser;

Xaa₈ is Ala, Ser or Thr;

Xaa₉ is Asp or Glu;

- Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
 Xaa₁₁ is Ala or Ser;
 Xaa₁₂ is Ala or Lys;
 Xaa₁₃ is Ala or Gln;
 5 Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
 Xaa₁₅ is Ala or Glu;
 Xaa₁₆ is Ala or Glu;
 Xaa₁₇ is Ala or Glu;
 Xaa₁₉ is Ala or Val;
 10 Xaa₂₀ is Ala or Arg;
 Xaa₂₁ is Ala or Leu;
 Xaa₂₂ is Ala, Phe, Tyr or naphthylalanine;
 Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine
 or Met;
 15 Xaa₂₄ is Ala, Glu or Asp;
 Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
 Xaa₂₆ is Ala or Leu;
 Xaa₂₇ is Ala or Lys;
 Xaa₂₈ is Ala or Asn;
 20 Z₁ is -OH,
 -NH₂
 Gly-Z₂,
 Gly Gly-Z₂,
 Gly Gly Xaa₃₁-Z₂,
 25 Gly Gly Xaa₃₁ Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,

Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,
Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂ or
Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂;
Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently Pro,
5 homoproline, 3Hyp, 4Hyp, thioproline,
N-alkylglycine, N-alkylpentylglycine or
N-alkylalanine; and
Z₂ is -OH or -NH₂;

provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈,
10 Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉,
Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈ are Ala.
Preferred N-alkyl groups for N-alkylglycine, N-
alkylpentylglycine and N-alkylalanine include lower alkyl
groups preferably of 1 to about 6 carbon atoms, more
15 preferably of 1 to 4 carbon atoms.

Preferred exendin agonist compounds include those
wherein Xaa₁ is His or Tyr. More preferably Xaa₁ is His.

Preferred are those compounds wherein Xaa₂ is Gly.

Preferred are those compounds wherein Xaa₁₄ is Leu,
20 pentylglycine or Met.

Preferred compounds are those wherein Xaa₂₅ is Trp or
Phe.

Preferred compounds are those where Xaa₆ is Phe or
naphthylalanine; Xaa₂₂ is Phe or naphthylalanine and
25 Xaa₂₃ is Ile or Val.

Preferred are compounds wherein Xaa₃₁, Xaa₃₆, Xaa₃₇ and
Xaa₃₈ are independently selected from Pro, homoproline,
thioproline and N-alkylalanine.

Preferably Z_1 is $-NH_2$.

Preferable Z_2 is $-NH_2$.

According to one aspect, preferred are compounds of formula (I) wherein Xaa_1 is His or Tyr, more preferably His; Xaa_2 is Gly; Xaa_6 is Phe or naphthylalanine; Xaa_{14} is Leu, pentylglycine or Met; Xaa_{22} is Phe or naphthylalanine; Xaa_{23} is Ile or Val; Xaa_{31} , Xaa_{36} , Xaa_{37} and Xaa_{38} are independently selected from Pro, homoproline, thioproline or N-alkylalanine. More preferably Z_1 is $-NH_2$.

According to an especially preferred aspect, especially preferred compounds include those of formula (I) wherein: Xaa_1 is His or Arg; Xaa_2 is Gly or Ala; Xaa_3 is Asp or Glu; Xaa_5 is Ala or Thr; Xaa_6 is Ala, Phe or naphthylalanine; Xaa_7 is Thr or Ser; Xaa_8 is Ala, Ser or Thr; Xaa_9 is Asp or Glu; Xaa_{10} is Ala, Leu or pentylglycine; Xaa_{11} is Ala or Ser; Xaa_{12} is Ala or Lys; Xaa_{13} is Ala or Gln; Xaa_{14} is Ala, Leu or pentylglycine; Xaa_{15} is Ala or Glu; Xaa_{16} is Ala or Glu; Xaa_{17} is Ala or Glu; Xaa_{19} is Ala or Val; Xaa_{20} is Ala or Arg; Xaa_{21} is Ala or Leu; Xaa_{22} is Phe or naphthylalanine; Xaa_{23} is Ile, Val or tert-butylglycine; Xaa_{24} is Ala, Glu or Asp; Xaa_{25} is Ala, Trp or Phe; Xaa_{26} is Ala or Leu; Xaa_{27} is Ala or Lys; Xaa_{28} is Ala or Asn; Z_1 is $-OH$, $-NH_2$, Gly- Z_2 , Gly Gly- Z_2 , Gly Gly Xaa_{31} - Z_2 , Gly Gly Xaa_{31} Ser- Z_2 , Gly Gly Xaa_{31} Ser Ser- Z_2 , Gly Gly Xaa_{31} Ser Ser Gly- Z_2 , Gly Gly Xaa_{31} Ser Ser Gly Ala- Z_2 , Gly Gly Xaa_{31} Ser Ser Gly Ala Xaa_{36} - Z_2 , Gly Gly Xaa_{31} Ser Ser Gly Ala Xaa_{36} Xaa_{37} - Z_2 , Gly Gly Xaa_{31} Ser Ser Gly Ala Xaa_{36} Xaa_{37} Xaa_{38} - Z_2 ; Xaa_{31} , Xaa_{36} , Xaa_{37} and Xaa_{38} being independently Pro homoproline, thioproline or N-

methyllalanine; and Z₂ being -OH or -NH₂; provided that no more than three of Xaa₃, Xaa₅, Xaa₆, Xaa₈, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆, Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈ are Ala. Especially
 5 preferred compounds include those having the amino acid sequence of SEQ. ID. NOS. 40-61.

According to an especially preferred aspect, provided are compounds where Xaa₁₄ is Leu, Ile, Val or pentylglycine, more preferably Leu or pentylglycine, and
 10 Xaa₂₅ is Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds will be less susceptible to oxidative degradation, both in vitro and in vivo, as well as during synthesis of the compound.

Exendin agonist compounds also include those
 15 described in U.S. Provisional Application No. 60/066,029, including compounds of the formula (III) [SEQ ID NO. 5]:

Xaa₁ Xaa₂ Xaa₃ Xaa₄ Xaa₅ Xaa₆ Xaa₇ Xaa₈ Xaa₉ Xaa₁₀
 Xaa₁₁ Xaa₁₂ Xaa₁₃ Xaa₁₄ Xaa₁₅ Xaa₁₆ Xaa₁₇ Ala Xaa₁₉ Xaa₂₀
 20 Xaa₂₁ Xaa₂₂ Xaa₂₃ Xaa₂₄ Xaa₂₅ Xaa₂₆ Xaa₂₇ Xaa₂₈-Z₁; wherein

Xaa₁ is His, Arg, Tyr, Ala, Norval, Val
 or Norleu;
 Xaa₂ is Ser, Gly, Ala or Thr;
 25 Xaa₃ is Ala, Asp or Glu;
 Xaa₄ is Ala, Norval, Val, Norleu or Gly;
 Xaa₅ is Ala or Thr;
 Xaa₆ is Phe, Tyr or naphthylalanine;

- Xaa₁ is Thr or Ser;
Xaa₂ is Ala, Ser or Thr;
Xaa₃ is Ala, Norval, Val, Norleu, Asp or Glu;
Xaa₁₀ is Ala, Leu, Ile, Val, pentylglycine or Met;
5 Xaa₁₁ is Ala or Ser;
Xaa₁₂ is Ala or Lys;
Xaa₁₃ is Ala or Gln;
Xaa₁₄ is Ala, Leu, Ile, pentylglycine, Val or Met;
Xaa₁₅ is Ala or Glu;
10 Xaa₁₆ is Ala or Glu;
Xaa₁₇ is Ala or Glu;
Xaa₁₉ is Ala or Val;
Xaa₂₀ is Ala or Arg;
Xaa₂₁ is Ala or Leu;
15 Xaa₂₂ is Phe, Tyr or naphthylalanine;
Xaa₂₃ is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met;
Xaa₂₄ is Ala, Glu or Asp;
Xaa₂₅ is Ala, Trp, Phe, Tyr or naphthylalanine;
20 Xaa₂₆ is Ala or Leu;
Xaa₂₇ is Ala or Lys;
Xaa₂₈ is Ala or Asn;
Z₁ is -OH,
-NH₂,
25 Gly-Z₂,
Gly Gly-Z₂,
Gly Gly Xaa₃₁-Z₂,
Gly Gly Xaa₃₁ Ser-Z₂,

Gly Gly Xaa₃₁ Ser Ser-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆-Z₂,
 5 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇-Z₂,
 Gly Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈-Z₂ or Gly
 Gly Xaa₃₁ Ser Ser Gly Ala Xaa₃₆ Xaa₃₇ Xaa₃₈ Xaa₃₉-Z₂; wherein
 Xaa₃₁, Xaa₃₆, Xaa₃₇ and Xaa₃₈ are independently
 Pro, homoproline, 3Hyp, 4Hyp, thioproline,
 10 N-alkylglycine, N-alkylpentylglycine or
 N-alkylalanine; and
 Z₂ is -OH or -NH₂;
 provided that no more than three of Xaa₃, Xaa₄, Xaa₅, Xaa₆,
 Xaa₈, Xaa₉, Xaa₁₀, Xaa₁₁, Xaa₁₂, Xaa₁₃, Xaa₁₄, Xaa₁₅, Xaa₁₆,
 15 Xaa₁₇, Xaa₁₉, Xaa₂₀, Xaa₂₁, Xaa₂₄, Xaa₂₅, Xaa₂₆, Xaa₂₇ and Xaa₂₈
 are Ala; and provided also that, if Xaa₁ is His, Arg or
 Tyr, then at least one of Xaa₃, Xaa₄ and Xaa₉ is Ala.

Definitions

20 In accordance with the present invention and as used
 herein, the following terms are defined to have the
 following meanings, unless explicitly stated otherwise.


The term "amino acid" refers to natural amino acids,
 unnatural amino acids, and amino acid analogs, all in
 25 their D and L stereoisomers if their structure allow such
 stereoisomeric forms. Natural amino acids include alanine
 (Ala), arginine (Arg), asparagine (Asn), aspartic acid
 (Asp), cysteine (Cys), glutamine (Gln), glutamic acid
 (Glu), glycine (Gly), histidine (His), isoleucine (Ile),

leucine (Leu), Lysine (Lys), methionine (Met),
phenylalanine (Phe), proline (Pro), serine (Ser),
threonine (Thr), typtophan (Trp), tyrosine (Tyr) and
valine (Val). Unnatural amino acids include, but are not
5 limited to azetidinecarboxylic acid, 2-aminoadipic acid,
3-aminoadipic acid, beta-alanine, aminopropionic acid, 2-
aminobutyric acid, 4-aminobutyric acid, 6-aminocaproic
acid, 2-aminoheptanoic acid, 2-aminoisobutyric acid, 3-
aminoisbutyric acid, 2-aminopimelic acid, tertiary-
10 butylglycine, 2,4-diaminoisobutyric acid, desmosine, 2,2'-
diaminopimelic acid, 2,3-diaminopropionic acid, N-
ethylglycine, N-ethylasparagine, homoproline,
hydroxylysine, allo-hydroxylysine, 3-hydroxyproline, 4-
hydroxyproline, isodesmosine, allo-isoleucine, N-
15 methylalanine, N-methylglycine, N-methylisoleucine, N-
methylpentylglycine, N-methylvaline, naphthalanine,
norvaline, norleucine, ornithine, pentylglycine, pipecolic
acid and thioproline. Amino acid analogs include the
natural and unnatural amino acids which are chemically
20 blocked, reversibly or irreversibly, or modified on their
N-terminal amino group or their side-chain groups, as for
example, methionine sulfoxide, methionine sulfone, S-
(carboxymethyl)-cysteine, S-(carboxymethyl)-cysteine
sulfoxide and S-(carboxymethyl)-cysteine sulfone.

25 The term "amino acid analog" refers to an amino acid
wherein either the C-terminal carboxy group, the N-
terminal amino group or side-chain functional group has
been chemically codified to another functional group. For
example, aspartic acid-(beta-methyl ester) is an amino

acid analog of aspartic acid; N-ethylglycine is an amino acid analog of glycine; or alanine carboxamide is an amino acid analog of alanine.

The term "amino acid residue" refers to radicals
5 having the structure: (1) $-C(O)-R-NH-$, wherein R typically is $-CH(R')$, wherein R' is an amino acid side chain, typically H or a carbon containing substituent; or (2),



10

wherein p is 1, 2 or 3 representing the azetidinecarboxylic acid, proline or pipercolic acid residues, respectively.

The term "lower" referred to herein in connection
15 with organic radicals such as alkyl groups defines such groups with up to and including about 6, preferably up to and including 4 and advantageously one or two carbon atoms. Such groups may be straight chain or branched chain.

20 "Pharmaceutically acceptable salt" includes salts of the compounds described herein derived from the combination of such compounds and an organic or inorganic acid. In practice the use of the salt form amounts to use of the base form. The compounds are useful in both free
25 base and salt form.

In addition, the following abbreviations stand for the following:

- "ACN" or "CH₃CN" refers to acetonitrile.
- "Boc", "tBoc" or "Tboc" refers to t-butoxy carbonyl.
- 5 "DCC" refers to N,N'-dicyclohexylcarbodiimide.
- "Fmoc" refers to fluorenylmethoxycarbonyl.
- "HBTU" refers to 2-(1H-benzotriazol-1-yl)-1,1,3,3,-tetramethyluronium hexafluorophosphate.
- "HOBt" refers to 1-hydroxybenzotriazole monohydrate.
- 10 "homoP" or hPro" refers to homoproline.
- "MeAla" or "Nme" refers to N-methylalanine.
- "naph" refers to naphthylalanine.
- "pG" or pGly" refers to pentylglycine.
- "tBuG" refers to tertiary-butylglycine.
- 15 "ThioP" or tPro" refers to thioproline.
- 3Hyp" refers to 3-hydroxyproline
- 4Hyp" refers to 4-hydroxyproline
- NAG" refers to N-alkylglycine
- NAPG" refers to N-alkylpentylglycine
- 20 "Norval" refers to norvaline
- "Norleu" refers to norleucine

Preparation of Compounds

- 25 The exendins and exendin agonists described herein may be prepared using standard solid-phase peptide synthesis techniques and preferably an automated or semiautomated peptide synthesizer. Typically, using such

techniques, an α -N-carbamoyl protected amino acid and an amino acid attached to the growing peptide chain on a resin are coupled at room temperature in an inert solvent such as dimethylformamide, N-methylpyrrolidinone or methylene chloride in the presence of coupling agents such as dicyclohexylcarbodiimide and 1-hydroxybenzotriazole in the presence of a base such as diisopropylethylamine. The α -N-carbamoyl protecting group is removed from the resulting peptide-resin using a reagent such as trifluoroacetic acid or piperidine, and the coupling reaction repeated with the next desired N-protected amino acid to be added to the peptide chain. Suitable N-protecting groups are well known in the art, with t-butyloxycarbonyl (tBoc) and fluorenylmethoxycarbonyl (Fmoc) being preferred herein.

The solvents, amino acid derivatives and 4-methylbenzhydryl-amine resin used in the peptide synthesizer may be purchased from Applied Biosystems Inc. (Foster City, CA). The following side-chain protected amino acids may be purchased from Applied Biosystems, Inc.: Boc-Arg(Mts), Fmoc-Arg(Pmc), Boc-Thr(Bzl), Fmoc-Thr(t-Bu), Boc-Ser(Bzl), Fmoc-Ser(t-Bu), Boc-Tyr(BrZ), Fmoc-Tyr(t-Bu), Boc-Lys(Cl-Z), Fmoc-Lys(Boc), Boc-Glu(Bzl), Fmoc-Glu(t-Bu), Fmoc-His(Trt), Fmoc-Asn(Trt), and Fmoc-Gln(Trt). Boc-His(BOM) may be purchased from Applied Biosystems, Inc. or Bachem Inc. (Torrance, CA). Anisole, dimethylsulfide, phenol, ethanedithiol, and thioanisole may be obtained from

Aldrich Chemical Company (Milwaukee, WI). Air Products and Chemicals (Allentown, PA) supplies HF. Ethyl ether, acetic acid and methanol may be purchased from Fisher Scientific (Pittsburgh, PA).

5 Solid phase peptide synthesis may be carried out with an automatic peptide synthesizer (Model 430A, Applied Biosystems Inc., Foster City, CA) using the NMP/HOBt (Option 1) system and tBoc or Fmoc chemistry (see, Applied Biosystems User's Manual for the ABI 430A Peptide
10 Synthesizer, Version 1.3B July 1, 1988, section 6, pp. 49-70, Applied Biosystems, Inc., Foster City, CA) with capping. Boc-peptide-resins may be cleaved with HF (-5° C to 0° C, 1 hour). The peptide may be extracted from the resin with alternating water and acetic acid, and the
15 filtrates lyophilized. The Fmoc-peptide resins may be cleaved according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc., 1990, pp. 6-12). Peptides may be also be assembled using an Advanced Chem Tech Synthesizer (Model MPS 350, Louisville,
20 Kentucky).

 Peptides may be purified by RP-HPLC (preparative and analytical) using a Waters Delta Prep 3000 system. A C4, C8 or C18 preparative column (10 μ , 2.2 x 25 cm; Vydac, Hesperia, CA) may be used to isolate peptides, and purity
25 may be determined using a C4, C8 or C18 analytical column (5 μ , 0.46 x 25 cm; Vydac). Solvents (A=0.1% TFA/water and B=0.1% TFA/CH₃CN) may be delivered to the analytical column at a flowrate of 1.0 ml/min and to the preparative column at 15 ml/min. Amino acid analyses may be performed

on the Waters Pico Tag system and processed using the Maxima program. Peptides may be hydrolyzed by vapor-phase acid hydrolysis (115° C, 20-24 h). Hydrolysates may be derivatized and analyzed by standard methods (Cohen, et al., The Pico Tag Method: A Manual of Advanced Techniques for Amino Acid Analysis, pp. 11-52, Millipore Corporation, Milford, MA (1989)). Fast atom bombardment analysis may be carried out by M-Scan, Incorporated (West Chester, PA).

Mass calibration may be performed using cesium iodide or cesium iodide/glycerol. Plasma desorption ionization analysis using time of flight detection may be carried out on an Applied Biosystems Bio-Ion 20 mass spectrometer. Electrospray mass spectroscopy may be carried out on a VG-Trio machine.

Peptide compounds useful in the invention may also be prepared using recombinant DNA techniques, using methods now known in the art. See, e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d Ed., Cold Spring Harbor (1989). Non-peptide compounds useful in the present invention may be prepared by art-known methods. For example, phosphate-containing amino acids and peptides containing such amino acids, may be prepared using methods known in the art. See, e.g., Bartlett and Landen, Biorg. Chem. 14:356-377 (1986).

The compounds described above are useful in view of their pharmacological properties. In particular, the compounds of the invention possess activity as agents to reduce food intake. They can be used to treat conditions or diseases which can be alleviated by reducing food

intake.

Compositions useful in the invention may conveniently be provided in the form of formulations suitable for parenteral (including intravenous, intramuscular and subcutaneous) or nasal or oral administration. In some cases, it will be convenient to provide an exendin or exendin agonist and another food-intake-reducing, plasma glucose-lowering or plasma lipid-lowering agent, such as amylin, an amylin agonist, a CCK, or a leptin, in a single composition or solution for administration together. In other cases, it may be more advantageous to administer the additional agent separately from said exendin or exendin agonist. A suitable administration format may best be determined by a medical practitioner for each patient individually. Suitable pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, e.g., Remington's Pharmaceutical Sciences by E.W. Martin. See also Wang, Y.J. and Hanson, M.A. "Parenteral Formulations of Proteins and Peptides: Stability and Stabilizers," Journal of Parenteral Science and Technology, Technical Report No. 10, Supp. 42:2S (1988).

Compounds useful in the invention can be provided as parenteral compositions for injection or infusion. They can, for example, be suspended in an inert oil, suitably a vegetable oil such as sesame, peanut, olive oil, or other acceptable carrier. Preferably, they are suspended in an aqueous carrier, for example, in an isotonic buffer solution at a pH of about 3.0 to 8.0, preferably at a pH of

about 3.5 to 5.0. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH buffering agents. Useful buffers include for example, sodium acetate/acetic acid buffers. A form of repository or "depot" slow release preparation may be used so that therapeutically effective amounts of the preparation are delivered into the bloodstream over many hours or days following transdermal injection or delivery.

The desired isotonicity may be accomplished using sodium chloride or other pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol, polyols (such as mannitol and sorbitol), or other inorganic or organic solutes. Sodium chloride is preferred particularly for buffers containing sodium ions.

The claimed compositions can also be formulated as pharmaceutically acceptable salts (e.g., acid addition salts) and/or complexes thereof. Pharmaceutically acceptable salts are non-toxic salts at the concentration at which they are administered. The preparation of such salts can facilitate the pharmacological use by altering the physical-chemical characteristics of the composition without preventing the composition from exerting its physiological effect. Examples of useful alterations in physical properties include lowering the melting point to facilitate transmucosal administration and increasing the solubility to facilitate the administration of higher

concentrations of the drug.

Pharmaceutically acceptable salts include acid addition salts such as those containing sulfate, hydrochloride, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, cyclohexylsulfamate and quinate. Pharmaceutically acceptable salts can be obtained from acids such as hydrochloric acid, sulfuric acid, phosphoric acid, sulfamic acid, acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, *p*-toluenesulfonic acid, cyclohexylsulfamic acid, and quinic acid. Such salts may be prepared by, for example, reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion on a suitable ion exchange resin.

Carriers or excipients can also be used to facilitate administration of the compound. Examples of carriers and excipients include calcium carbonate, calcium phosphate, various sugars such as lactose, glucose, or sucrose, or types of starch, cellulose derivatives, gelatin, vegetable oils, polyethylene glycols and physiologically compatible solvents. The compositions or pharmaceutical composition can be administered by different routes including intravenously, intraperitoneal, subcutaneous, and intramuscular, orally, topically, transmucosally, or by

pulmonary inhalation.

If desired, solutions of the above compositions may be thickened with a thickening agent such as methyl cellulose.

They may be prepared in emulsified form, either water in oil or oil in water. Any of a wide variety of pharmaceutically acceptable emulsifying agents may be employed including, for example, acacia powder, a non-ionic surfactant (such as a Tween), or an ionic surfactant (such as alkali polyether alcohol sulfates or sulfonates, e.g., a Triton).

Compositions useful in the invention are prepared by mixing the ingredients following generally accepted procedures. For example, the selected components may be simply mixed in a blender or other standard device to produce a concentrated mixture which may then be adjusted to the final concentration and viscosity by the addition of water or thickening agent and possibly a buffer to control pH or an additional solute to control tonicity.

For use by the physician, the compositions will be provided in dosage unit form containing an amount of an exendin or exendin agonist, for example, exendin-3, and/or exendin-4, with or without another food intake-reducing, plasma glucose-lowering or plasma lipid-lowering agent. Therapeutically effective amounts of an exendin or exendin agonist for use in reducing food intake are those that suppress appetite at a desired level. As will be recognized by those in the field, an effective amount of therapeutic agent will vary with many factors including the age and weight of the patient, the patient's physical

condition, the blood sugar level and other factors.

The effective daily appetite-suppressing dose of the compounds will typically be in the range of about 10 to 30 μ g to about 5 mg/day, preferably about 10 to 30 μ g to about 2 mg/day and more preferably about 10 to 100 μ g to about 1 mg/day, most preferably about 30 μ g to about 500 μ g/day, for a 70 kg patient, administered in a single or divided doses. The exact dose to be administered is determined by the attending clinician and is dependent upon where the particular compound lies within the above quoted range, as well as upon the age, weight and condition of the individual. Administration should begin whenever the suppression of food intake, or weight lowering is desired, for example, at the first sign of symptoms or shortly after diagnosis of obesity, diabetes mellitus, or insulin-resistance syndrome. Administration may be by injection, preferably subcutaneous or intramuscular. Orally active compounds may be taken orally, however dosages should be increased 5-10 fold.

The optimal formulation and mode of administration of compounds of the present application to a patient depend on factors known in the art such as the particular disease or disorder, the desired effect, and the type of patient. While the compounds will typically be used to treat human subjects they may also be used to treat similar or identical diseases in other vertebrates such as other primates, farm animals such as swine, cattle and poultry, and sports animals and pets such as horses, dogs and cats.

To assist in understanding the present invention, the following Examples are included. The experiments relating to this invention should not, of course, be construed as specifically limiting the invention and such variations of the invention, now known or later developed, which would be within the purview of one skilled in the art are considered to fall within the scope of the invention as described herein and hereinafter claimed.

EXAMPLE 1: Exendin Injections Reduced the Food Intake of
Normal Mice

All mice (NIH:Swiss mice) were housed in a stable environment of 22 (\pm 2) $^{\circ}$ C, 60 (\pm 10) % humidity and a 12:12 light:dark cycle; with lights on at 0600. Mice were housed in groups of four in standard cages with *ad libitum* access to food (Teklad: LM 485; Madison, WI) and water except as noted, for at least two weeks before the experiments.

All experiments were conducted between the hours of 0700 and 0900. The mice were food deprived (food removed at 1600 hr from all animals on day prior to experiment) and individually housed. All mice received an intraperitoneal injection (5 μ l/kg) of either saline or exendin-4 at doses of 0.1, 1.0, 10 and 100 μ g/kg and were immediately presented with a pre-weighed food pellet (Teklad LM 485). The food pellet was weighed at 30-minute, 1-hr, 2-hr and 6-hr intervals to determine the amount of food eaten.

Figure 1 depicts cumulative food intake over periods of 0.5, 1, 2 and 6hr in overnight-fasted normal NIH:Swiss

mice following ip injection of saline, 2 doses of GLP-1, or 4 doses of exendin-4. At doses up to 100µg/kg, GLP-1 had no effect on food intake measured over any period, a result consistent with that previously reported (Bhavsar, S.P., et al., Soc. Neurosci. Abstr. 21:460 (188.8) (1995); and 5 Turton, M.D., Nature, 379:69-72, (1996)).

In contrast, exendin-4 injections potently and dose-dependently inhibited food intake. The ED₅₀ for inhibition of food intake over 30 min was 1µg/kg, which is a level 10 about as potent as amylin (ED₅₀ 3.6µg/kg) or the prototypical peripheral satiety agent, CCK (ED₅₀ 0.97µg/kg) as measured in this preparation. However, in contrast to the effects of amylin or CCK, which abate after 1-2 hours, the inhibition of food intake with exendin-4 was still 15 present after at least 6 hours after injection.

EXAMPLE 2: Exendin Reduced the Food Intake of Obese Mice

All mice (female ob/ob mice) were housed in a stable environment of 22 (±2)° C, 60 (±10) % humidity and a 12:12 20 light:dark cycle; with lights on at 0600. Mice were housed in groups of four in standard cages with *ad libitum* access to food (Teklad: LM 485) and water except as noted, for at least two weeks before the experiments.

All experiments were conducted between the hours of 25 0700 and 0900. The mice were food deprived (food removed at 1600 hr from all animals on day prior to experiment) and individually housed. All mice received an intraperitoneal injection (5 µl/kg) of either saline or exendin-4 at doses

of 0.1, 1.0 and 10 $\mu\text{g/kg}$ (female ob/ob mice) and were immediately presented with a pre-weighed food pellet (Teklad LM 485). The food pellet was weighed at 30-minute, 1 -hr, 2-hr and 6-hr intervals to determine the amount of food eaten.

Figure 2 depicts the effect of exendin-4 in the ob/ob mouse model of obesity. The obese mice had a similar food intake-related response to exendin as the normal mice. Moreover, the obese mice were not hypersensitive to exendin, as has been observed with amylin and leptin (Young, A.A., et al., Program and Abstracts, 10th International Congress of Endocrinology, June 12-15, 1996 San Francisco, pg 419 (P2-58)).

EXAMPLE 3: Intracerebroventricular Injections of Exendin
Inhibited Food Intake in Rats

All rats (Harlan Sprague-Dawley) were housed in a stable environment of 22 (± 2)° C, 60 (± 10)% humidity and a 12:12 light:dark cycle; with lights on at 0600. Rats were obtained from Zivic Miller with an intracerebroventricular cannula (ICV cannula) implanted (coordinates determined by actual weight of animals and referenced to Paxinos, G. and Watson, C. "The Rat Brain in stereotaxic coordinates," second edition. Academic Press) and were individually housed in standard cages with *ad libitum* access to food (Teklad: LM 485) and water for at least one week before the experiments.

All injections were given between the hours of 1700

and 1800. The rats were habituated to the ICV injection procedure at least once before the ICV administration of compound. All rats received an ICV injection (2 μ l/30 seconds) of either saline or exendin-4 at doses of 0.01, 0.03, 0.1, 0.3, and 1.0 μ g. All animals were then presented with pre-weighed food (Teklad LM 485) at 1800, when the lights were turned off. The amount of food left was weighed at 2-hr, 12-hr and 24-hr intervals to determine the amount of food eaten by each animal.

Figure 3 depicts a dose-dependent inhibition of food intake in rats that received doses greater than 0.1 μ g/rat. The ED₅₀ was \approx 0.1 μ g, exendin-4 is thus \approx 100-fold more potent than intracerebroventricular injections of GLP-1 as reported by Turton, M.D., et al. (Nature 379:69-72 (1996)).

EXAMPLE 4: Exendin Agonists Reduced the Food Intake in Mice

All mice (NIH:Swiss mice) were housed in a stable environment of 22 (\pm 2)° C, 60 (\pm 10) % humidity and a 12:12 light:dark cycle; with lights on at 0600. Mice were housed in groups of four in standard cages with *ad libitum* access to food (Teklad: LM 485; Madison, WI) and water except as noted, for at least two weeks before the experiments.

All experiments were conducted between the hours of 0700 and 0900. The mice were food deprived (food removed at 1600 hr from all animals on day prior to experiment) and individually housed. All mice received an

intraperitoneal injection (5 μ l/kg) of either saline or test compound at doses of 1, 10, and 100 μ g/kg and immediately presented with a food pellet (Teklad LM 485).

5 The food pellet was weighed at 30-minute, 1-hr, 2-hr and 6-hr intervals to determine the amount of food eaten.

Figure 4 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or exendin-4 (1-30) ("Compound 1") in doses of 1, 10 and 100
10 μ g/kg.

Figure 5 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or exendin-4 (1-30) amide ("Compound 2") in doses of 1, 10
15 and 100 μ g/kg.

Figure 6 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or exendin-4 (1-28) amide ("Compound 3") in doses of 1, 10
20 and 100 μ g/kg.

Figure 7 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or ¹⁴Leu, ²⁵Phe exendin-4 amide ("Compound 4") in doses of 1, 10
25 and 100 μ g/kg.

Figure 8 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or ¹⁴Leu, ²⁵Phe exendin-4 (1-28) amide ("Compound 5") in doses

of 1, 10 and 100 $\mu\text{g/kg}$.

Figure 9 depicts the cumulative food intake over periods of 0.5, 1, 2 and 6 hr in overnight-fasted normal NIH:Swiss mice following ip injection of saline or
5 ^{14}Leu , ^{22}Ala , ^{25}Phe exendin-4 (1-28) amide ("Compound 6") in doses of 1, 10 and 100 $\mu\text{g/kg}$.

EXAMPLE 5

Preparation of amidated peptide having SEQ. ID. NO. 9

10

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.). In
15 general, single-coupling cycles were used throughout the synthesis and Fast Moc (HBTU activation) chemistry was employed. However, at some positions coupling was less efficient than expected and double couplings were required. In particular, residues Asp_9 , Thr , and Phe_6 all
20 required double coupling. Deprotection (Fmoc group removal) of the growing peptide chain using piperidine was not always efficient. Double deprotection was required at positions Arg_{20} , Val_{19} and Leu_{14} . Final deprotection of the completed peptide resin was achieved using a mixture of
25 triethylsilane (0.2 mL), ethanedithiol (0.2 mL), anisole (0.2 mL), water (0.2 mL) and trifluoroacetic acid (15 mL) according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc.) The peptide was precipitated in ether/water (50 mL) and centrifuged. The

precipitate was reconstituted in glacial acetic acid and lyophilized. The lyophilized peptide was dissolved in water). Crude purity was about 55%.

Used in purification steps and analysis were Solvent
5 A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

The solution containing peptide was applied to a preparative C-18 column and purified (10% to 40% Solvent B in Solvent A over 40 minutes). Purity of fractions was determined isocratically using a C-18 analytical column.
10 Pure fractions were pooled furnishing the above-identified peptide. Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.5 minutes. Electrospray Mass Spectrometry (M):
15 calculated 4131.7; found 4129.3.

EXAMPLE 6

Preparation of Peptide having SEQ. ID. NO. 10

20 The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.),
25 cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 25% to 75% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 21.5

minutes. Electrospray Mass Spectrometry (M): calculated 4168.6; found 4171.2.

EXAMPLE 7

5 Preparation of Peptide having SEO. ID. NO. 11

The above-identified peptide was assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
10 Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis were Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
15 Solvent A over 30 minutes) of the lyophilized peptide gave
product peptide having an observed retention time of 17.9
minutes. Electrospray Mass Spectrometry (M): calculated
4147.6; found 4150.2.

20 EXAMPLE 8

Preparation of Peptide having SEO. ID. NO. 12

The above-identified peptide was assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
25 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis were Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

Analytical RP-HPLC (gradient 35% to 65% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 19.7 minutes. Electrospray Mass Spectrometry (M): calculated
5 4212.6; found 4213.2.

EXAMPLE 9

Preparation of Peptide having SEQ. ID. NO. 13

10 The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.),
15 cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 50% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave
20 product peptide having an observed retention time of 16.3 minutes. Electrospray Mass Spectrometry (M): calculated 4262.7; found 4262.4.

EXAMPLE 10

Preparation of Peptide having SEQ. ID. NO. 14

25 The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.),

cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4172.6

10

EXAMPLE 11Preparation of Peptide having SEQ. ID. NO. 15

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4224.7.

25

EXAMPLE 12Preparation of Peptide having SEQ. ID. NO. 16

5 The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
10 similar way to Example 5. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
15 product peptide. Electrospray Mass Spectrometry (M):
calculated 4172.6

EXAMPLE 13Preparation of Peptide having SEQ. ID. NO. 17

20 The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
25 cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is

then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4186.6

5

EXAMPLE 14Preparation of Peptide having SEQ. ID. NO. 18

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
10 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
15 Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4200.7

20

EXAMPLE 15Preparation of Peptide having SEQ. ID. NO. 19

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
25 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A

(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
5 product peptide. Electrospray Mass Spectrometry (M):
calculated 4200.7

EXAMPLE 16

Preparation of Peptide having SEQ. ID. NO. 20

10

The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
15 cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
20 then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 4202.7.

EXAMPLE 17

25

Preparation of Peptide having SEQ. ID. NO. 21

The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using

Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
5 Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 4145.6.

10

EXAMPLE 18Preparation of Peptide having SEQ. ID. NO. 22

The above-identified peptide is assembled on 4-(2'-
15 4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis are Solvent A
20 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
25 calculated 4184.6.

EXAMPLE 19Preparation of Peptide having SEQ. ID. NO. 23

5 The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
10 similar way to Example 5. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
15 product peptide. Electrospray Mass Spectrometry (M):
calculated 4145.6.

EXAMPLE 20Preparation of Peptide having SEQ. ID. NO. 24

20 The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
25 Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in

Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4224.7.

5

EXAMPLE 21

Preparation of Peptide having SEQ. ID. NO. 25

The above-identified peptide is assembled on 4-(2'-
10 4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis are Solvent A
15 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
20 calculated 4172.6.

EXAMPLE 22

Preparation of Peptide having SEQ. ID. NO. 26

25 The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a

similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4115.5.

EXAMPLE 23

Preparation of Peptide having SEQ. ID. NO. 27

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4188.6.

EXAMPLE 24Preparation of Peptide having SEQ. ID. NO. 28

5 The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis are Solvent A
10 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
15 calculated 4131.6.

EXAMPLE 25Preparation of Peptide having SEQ. ID. NO. 29

20 The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis are Solvent A
25 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the

product peptide. Electrospray Mass Spectrometry (M):
calculated 4172.6.

EXAMPLE 26

5 Preparation of Peptide having SEQ. ID. NO. 30

The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
10 Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
15 Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 4145.6.

EXAMPLE 27

20 Preparation of Peptide having SEQ. ID. NO. 31

The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
25 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Additional double couplings are
required at the thioproline positions 38, 37, 36 and 31.

Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4266.8.

EXAMPLE 28

Preparation of Peptide having SEQ. ID. NO. 32

10

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Additional double couplings are required at the thioproline positions 38, 37 and 36. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4246.8.

25

EXAMPLE 29

Preparation of Peptide having SEQ. ID. NO. 33

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide

norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Additional double couplings are required at the homoproline positions 38, 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4250.8.

EXAMPLE 30

Preparation of Peptide having SEQ. ID. NO. 34

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Additional double couplings are required at the homoproline positions 38, 37, and 36. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4234.8.

EXAMPLE 31Preparation of Peptide having SEQ. ID. NO. 35

5 The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
10 similar way to Example 5. Additional double couplings are
required at the thioproline positions 38, 37, 36 and 31.
Used in analysis are Solvent A (0.1% TFA in water) and
Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
30% to 60% Solvent B in Solvent A over 30 minutes) of the
15 lyophilized peptide is then carried out to determine the
retention time of the product peptide. Electrospray Mass
Spectrometry (M): calculated 4209.8.

EXAMPLE 32

20 Preparation of Peptide having SEQ. ID. NO. 36

 The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
25 Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Additional double couplings are
required at the homoproline positions 38, 37, 36 and 31.
Used in analysis are Solvent A (0.1% TFA in water) and

Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4193.7.

EXAMPLE 33

Preparation of Peptide having SEQ. ID. NO. 37

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 5. Additional double couplings are required at the N-methylalanine positions 38, 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3858.2.

EXAMPLE 34

Preparation of Peptide having SEQ. ID. NO. 38

The above-identified peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using

Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 5. Additional double couplings are
required at the N-methylalanine positions 38, 37 and 36.
5 Used in analysis are Solvent A (0.1% TFA in water) and
Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
30% to 60% Solvent B in Solvent A over 30 minutes) of the
lyophilized peptide is then carried out to determine the
retention time of the product peptide. Electrospray Mass
10 Spectrometry (M): calculated 3940.3.

EXAMPLE 35

Preparation of Peptide having SEQ. ID. NO. 39

15

The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
20 cleaved from the resin, deprotected and purified in a
similar way to Example 5. Additional double couplings are
required at the N-methylalanine positions 38, 37, 36 and
31. Used in analysis are Solvent A (0.1% TFA in water) and
Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient
25 30% to 60% Solvent B in Solvent A over 30 minutes) of the
lyophilized peptide is then carried out to determine the
retention time of the product peptide. Electrospray Mass
Spectrometry (M): calculated 3801.1.

EXAMPLE 36Preparation of C-terminal carboxylic acid Peptides
corresponding to the above C-terminal amide sequences.

5

The above peptides of Examples 5 to 35 are assembled on the so called Wang resin (p-alkoxybenzylalcohol resin (Bachem, 0.54 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin,
10 deprotected and purified in a similar way to Example 5. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the
15 retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

EXAMPLE 37

20

Preparation of Peptide having SEQ ID NO. 7

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly-NH₂ [SEQ. ID. NO. 7]

25

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.). In
30 general, single-coupling cycles were used throughout the

synthesis and Fast Moc (HBTU activation) chemistry was employed. Deprotection (Fmoc group removal) of the growing peptide chain was achieved using piperidine. Final deprotection of the completed peptide resin was achieved using a mixture of triethylsilane (0.2 mL), ethanedithiol (0.2 mL), anisole (0.2 mL), water (0.2 mL) and trifluoroacetic acid (15 mL) according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc.) The peptide was precipitated in ether/water (50 mL) and centrifuged. The precipitate was reconstituted in glacial acetic acid and lyophilized. The lyophilized peptide was dissolved in water). Crude purity was about 75%.

Used in purification steps and analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). The solution containing peptide was applied to a preparative C-18 column and purified (10% to 40% Solvent B in Solvent A over 40 minutes). Purity of fractions was determined isocratically using a C-18 analytical column. Pure fractions were pooled furnishing the above-identified peptide. Analytical RP-HPLC (gradient 30% to 50% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 18.9 minutes. Electrospray Mass Spectrometry (M): calculated 3408.0; found 3408.9.

EXAMPLE 38Preparation of Peptide having SEQ ID NO. 40

5 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 40]

The above amidated peptide was assembled on 4-(2'-4'-
10 dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 37. Used in analysis were Solvent A
15 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 40% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide gave
product peptide having an observed retention time of 17.9
minutes. Electrospray Mass Spectrometry (M): calculated
20 3294.7; found 3294.8.

EXAMPLE 39Preparation of Peptide having SEQ ID NO. 41

25

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 41]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 29% to 36% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 20.7 minutes. Electrospray Mass Spectrometry (M): calculated 3237.6; found 3240.

EXAMPLE 40

Preparation of Peptide having SEQ ID NO. 42

His Ala Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 42]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 15.2 minutes. Electrospray Mass Spectrometry (M): calculated
5 3251.6; found 3251.5.

EXAMPLE 41

Preparation of Peptide having SEQ ID NO. 43

10 His Gly Glu Gly Ala Phe Thr Ser Asp Leu Ser Lys Gln
Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys
Asn-NH₂ [SEQ. ID. NO. 43]

The above amidated peptide was assembled on 4-(2'-4'-
15 dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 37. Used in analysis were Solvent A
20 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 36% to 46% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide gave
product peptide having an observed retention time of 13.1
minutes. Electrospray Mass Spectrometry (M): calculated
25 3207.6; found 3208.3.

EXAMPLE 42Preparation of Peptide having SEQ ID NO: 44

His Gly Glu Gly Thr Ala Thr Ser Asp Leu Ser Lys Gln Leu
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 44]

The above amidated peptide was assembled on 4-(2'-4'-
dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
10 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 37. Used in analysis were Solvent
A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
15 Analytical RP-HPLC (gradient 35% to 45% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide gave
product peptide having an observed retention time of 12.8
minutes. Electrospray Mass Spectrometry (M): calculated
3161.5; found 3163.

20

EXAMPLE 43Preparation of Peptide having SEQ ID NO. 45

His Gly Glu Gly Thr Phe Thr Ala Asp Leu Ser Lys Gln Leu
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 45]

The above-identified amidated peptide was assembled

on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
5 purified in a similar way to Example 37. Used in analysis
were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
gave product peptide having an observed retention time of
10 15.2 minutes. Electrospray Mass Spectrometry (M):
calculated 3221.6; found 3222.7.

EXAMPLE 44

Preparation of Peptide having SEQ ID NO. 46

15

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 46]

20

The above-identified amidated peptide was assembled
on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
25 purified in a similar way to Example 37. Used in analysis
were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 34% to 44% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide

gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3199.4.

5

EXAMPLE 45Preparation of Peptide having SEQ ID NO. 47

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ala Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 47]

10

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 15.7 minutes. Electrospray Mass Spectrometry (M): calculated 3221.6; found 3221.6.

15

20

25

EXAMPLE 46Preparation of Peptide having SEQ ID NO. 48

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Ala Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 48]

30

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 18.1 minutes. Electrospray Mass Spectrometry (M): calculated 3180.5; found 3180.9.

15

EXAMPLE 47Preparation of Peptide having SEQ ID NO. 49

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Ala Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 49]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Compound 1. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

gave product peptide having an observed retention time of 17.0 minutes. Electrospray Mass Spectrometry (M): calculated 3180.6; found 3182.8.

5

EXAMPLE 48Preparation of Peptide having SEQ ID NO. 50

10

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Ala
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 50]

15

20

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.9 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3195.9.

25

EXAMPLE 49Preparation of Peptide having SEQ ID NO. 51

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu

Ala Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 51]

5 The above-identified amidated peptide was assembled
on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
10 purified in a similar way to Example 37. Used in analysis
were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
gave product peptide having an observed retention time of
17.9 minutes. Electrospray Mass Spectrometry (M):
15 calculated 3179.6; found 3179.0.

EXAMPLE 50

Preparation of Peptide having SEQ ID NO. 52

20

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Ala Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 52]

25

The above-identified amidated peptide was assembled
on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and

purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.0.

EXAMPLE 51

Preparation of Peptide having SEQ ID NO. 53

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Ala Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 53]

The above-identified peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 13.7 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3179.0.

EXAMPLE 52Preparation of Peptide having SEQ ID NO. 54

5 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Ala Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 54]

The above-identified amidated peptide was assembled
10 on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
15 were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 35% to 45% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
gave product peptide having an observed retention time of
14.0 minutes. Electrospray Mass Spectrometry (M):
20 calculated 3209.6; found 3212.8.

EXAMPLE 53Preparation of Peptide having SEQ ID NO. 55

25 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Ala Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 55]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.3 minutes. Electrospray Mass Spectrometry (M): calculated 3152.5; found 3153.5.

EXAMPLE 54

Preparation of Peptide having SEQ ID NO. 56

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Ala Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 56]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 35% to 45% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

gave product peptide having an observed retention time of 12.1 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3197.7.

5

EXAMPLE 55Preparation of Peptide having SEQ ID NO. 57

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Ala Phe Leu Lys Asn-NH₂
10 [SEQ. ID. NO. 57]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B
20 in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 10.9 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.5.

25

EXAMPLE 56Preparation of Peptide having SEQ ID NO. 58

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Ala Leu Lys Asn-NH₂

[SEQ. ID. NO. 58]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 17.5 minutes. Electrospray Mass Spectrometry (M): calculated 3161.5; found 3163.0.

15

EXAMPLE 57

Preparation of Peptide having SEQ ID NO. 59

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Ala Lys Asn-NH₂
[SEQ. ID. NO. 59]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 32% to 42% Solvent B

in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 19.5 minutes. Electrospray Mass Spectrometry (M): calculated 3195.5; found 3199.

5

EXAMPLE 58Preparation of Peptide having SEQ ID NO. 60

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Ala Asn-NH₂
[SEQ. ID. NO. 60]

The above-identified amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
15 acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis
20 were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 38% to 48% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.5 minutes. Electrospray Mass Spectrometry (M): calculated 3180.5; found 3183.7.

25

EXAMPLE 59Preparation of Peptide having SEQ ID NO. 61

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Ala-NH₂
[SEQ. ID. NO. 61]

The above-identified amidated peptide was assembled
on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
15 were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 34% to 44% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
gave product peptide having an observed retention time of
22.8 minutes. Electrospray Mass Spectrometry (M):
calculated 3194.6; found 3197.6.

20

EXAMPLE 60Preparation of Peptide having SEQ ID NO. 62

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH₂ [SEQ. ID. NO.
62]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4099.6.

EXAMPLE 61

Preparation of Peptide having SEQ ID NO. 63

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH₂ [SEQ. ID. NO.
63]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
5 calculated 4042.5.

EXAMPLE 62

Preparation of Peptide having SEQ ID NO: 64

10 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro-NH₂ [SEQ. ID. NO. 64]

The above-identified peptide is assembled on 4-(2'-
15 4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 37. Used in analysis are Solvent A
20 (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
25 calculated 4002.4

EXAMPLE 63Preparation of Peptide having SEQ ID NO: 65

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro-NH₂ [SEQ. ID. NO. 65]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
20 calculated 3945.4.

EXAMPLE 64Preparation of Peptide having SEQ ID NO. 66

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln
25 Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys
Asn Gly Gly Pro Ser Ser Gly Ala Pro-NH₂ [SEQ. ID. NO. 66]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
5 purified in a similar way to Example 37. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
10 product peptide. Electrospray Mass Spectrometry (M):
calculated 3905.3.

EXAMPLE 65

Preparation of Peptide having SEQ ID NO. 67

15

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro-NH₂ [SEQ. ID. NO. 67]

20

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
25 purified in a similar way to Example 37. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3848.2.

5

EXAMPLE 66Preparation of Peptide having SEQ ID NO. 68

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
10 Gly Gly Pro Ser Ser Gly Ala-NH₂, [SEQ. ID. NO. 68]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
20 in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3808.2.

25

EXAMPLE 67Preparation of Peptide having SEQ ID NO. 69

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 69]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3751.1.

20

EXAMPLE 68Preparation of Peptide having SEQ ID NO. 70

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly-NH₂ [SEQ. ID. NO. 70]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
5 purified in a similar way to Example 37. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
10 product peptide. Electrospray Mass Spectrometry (M):
calculated 3737.1.

EXAMPLE 69

Preparation of Peptide having SEQ ID NO. 71

15

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser Gly-NH₂ [SEQ. ID. NO. 71]

20

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
25 purified in a similar way to Example 37. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3680.1.

5

EXAMPLE 70Preparation of Peptide having SEQ ID NO. 72

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
10 Gly Gly Pro Ser Ser-NH₂ [SEQ. ID. NO. 72]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
20 in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3680.1

25

EXAMPLE 71Preparation of Peptide having SEQ ID NO. 73

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
30 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser-NH₂ [SEQ. ID. NO. 73]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3623.0.

15

EXAMPLE 72Preparation of Peptide having SEQ ID NO. 74

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
20 Gly Gly Pro Ser-NH₂ [SEQ. ID. NO. 74]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
5 calculated 3593.0

EXAMPLE 73

Preparation of Peptide having SEQ ID NO. 75

10 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser-NH₂ [SEQ. ID. NO. 75]

The above-identified amidated peptide is assembled on
15 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
20 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
25 calculated 3535.9

EXAMPLE 74Preparation of Peptide having SEQ ID NO. 76

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro-NH₂ [SEQ. ID. NO. 76]

The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
10 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 37. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
15 Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3505.9.

EXAMPLE 75Preparation of Peptide having SEQ ID NO. 77

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro-NH₂ [SEQ. ID. NO. 77]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
5 purified in a similar way to Example 37. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
10 product peptide. Electrospray Mass Spectrometry (M):
calculated 3448.8.

EXAMPLE 76

Preparation of Peptide having SEQ ID NO. 78

15

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly-NH₂ [SEQ. ID. NO. 78]

20

The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
25 similar way to Example 37. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is

then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3351.7.

5

EXAMPLE 77Preparation of Peptide having SEQ ID NO. 79

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly-NH₂ [SEQ. ID. NO. 79]

The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
15 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 37. Used in analysis are Solvent A
(0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
20 Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3351.8.

25

EXAMPLE 78Preparation of Peptide having SEQ ID NO. 80

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly-NH₂ [SEQ. ID. NO. 80]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3294.7.

20

EXAMPLE 79Preparation of Peptide having SEQ ID NO. 81

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly tPro Ser Ser Gly Ala tPro tPro tPro-NH₂ [SEQ. ID.
NO. 81]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Double couplings are required at residues 37,36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4197.1.

15

EXAMPLE 80Preparation of Peptide having SEQ ID NO. 82

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
20 Gly Gly Pro Ser Ser Gly Ala tPro tPro tPro-NH₂ [SEQ. ID.
NO. 82]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Double couplings

are required at residues 37, 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4179.1.

EXAMPLE 81

Preparation of Peptide having SEQ ID NO. 83

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly NMeala Ser Ser Gly Ala Pro Pro-NH₂ [SEQ. ID. NO.
83]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Double couplings are required at residues 36 and 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3948.3.

EXAMPLE 82

Preparation of Peptide having SEQ ID NO. 84

5

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly NMeala Ser Ser Gly Ala NMeala Nmeala-NH₂ [SEQ. ID.
NO. 84]

10

15

20

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Double couplings
are required at residues 36 and 31. Used in analysis are
Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in
ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3840.1.

EXAMPLE 83Preparation of Peptide having SEQ ID NO. 85

5 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly hPro Ser Ser Gly Ala hPro hPro-NH₂ [SEQ. ID. NO.
85]

10 The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
15 purified in a similar way to Example 37. Double couplings
are required at residues 36 and 31. Used in analysis are
Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in
ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
20 product peptide. Electrospray Mass Spectrometry (M):
calculated 4050.1.

EXAMPLE 84Preparation of Peptide having SEQ ID NO. 86

25 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly hPro Ser Ser Gly Ala hPro-NH₂ [SEQ. ID. NO. 86]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. A double coupling is required at residue 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3937.1

15

EXAMPLE 85

Preparation of Peptide having SEQ ID NO. 87

Arg Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 87]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis

are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
5 calculated 3827.2.

EXAMPLE 86

Preparation of Peptide having SEQ ID NO. 88

10

His Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly-NH₂ [SEQ. ID. NO. 88]

15

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and
20 purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the
25 product peptide. Electrospray Mass Spectrometry (M): calculated 3394.8.

EXAMPLE 87Preparation of Peptide having SEQ ID NO. 89

His Gly Glu Gly Thr Naphthylala Thr Ser Asp Leu Ser Lys
5 Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu
Lys Asn-NH₂ [SEQ. ID. NO. 89]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3289.5.

20

EXAMPLE 88Preparation of Peptide having SEQ ID NO. 90

His Gly Glu Gly Thr Phe Ser Ser Asp Leu Ser Lys Gln Met
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 90]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
5 purified in a similar way to Example 37. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
10 product peptide. Electrospray Mass Spectrometry (M):
calculated 3280.7.

EXAMPLE 89

Preparation of Peptide having SEQ ID NO. 91

15

His Gly Glu Gly Thr Phe Ser Thr Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 91]

20

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
25 purified in a similar way to Example 37. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3294.7.

5

EXAMPLE 90Preparation of Peptide having SEQ ID NO. 92

His Gly Glu Gly Thr Phe Thr Ser Glu Leu Ser Lys Gln Met
Ala Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 92]

10

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3250.7.

15

20

EXAMPLE 91

25

Preparation of Peptide having SEQ ID NO. 93

His Gly Glu Gly Thr Phe Thr Ser Asp pentylgly Ser Lys Gln
Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys

Asn-NH₂ [SEQ. ID. NO. 93]

5 The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
10 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3253.5.

15

EXAMPLE 92

Preparation of Peptide having SEQ ID NO. 94

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
20 Glu Glu Glu Ala Val Arg Leu Naphthylala Ile Glu Phe Leu
Lys Asn-NH₂ [SEQ. ID. NO. 94]

25 The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis

are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
5 calculated 3289.5.

EXAMPLE 93

Preparation of Peptide having SEQ ID NO. 95

10

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe tButylgly Glu Trp Leu Lys
Asn-NH₂ [SEQ. ID. NO. 95]

15

20

25

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
calculated 3183.4.

EXAMPLE 94Preparation of Peptide having SEQ ID NO. 96

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Asp Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 96]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3237.6.

20

EXAMPLE 95Preparation of Peptide having SEQ ID NO. 97

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser-NH₂ [SEQ. ID. NO. 97]

The above-identified amidated peptide is assembled on

4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
5 purified in a similar way to Example 37. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
10 product peptide. Electrospray Mass Spectrometry (M):
calculated 3637.9.

EXAMPLE 96

Preparation of Peptide having SEQ ID NO. 98

15

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly-NH₂ [SEQ. ID. NO. 98]

20

25

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3309.7.

5

EXAMPLE 97Preparation of Peptide having SEO ID NO. 99

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
10 Gly Gly hPro Ser Ser Gly Ala hPro hPro-NH₂ [SEQ. ID. NO.
99]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
15 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 37. Double couplings
are required at residues 36 and 31. Used in analysis are
20 Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in
ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is
then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
25 calculated 3711.1.

EXAMPLE 98

Preparation of C-terminal carboxylic acid peptides
corresponding to the above C-terminal amide sequences for
SEQ ID NOS. 7, 40-61, 68-75, 78-80 and 87-96

5

Peptides having the sequences of SEQ ID NOS. 7, 40-61, 68-75, 78-80 and 87-96 are assembled on the so called Wang resin (p-alkoxybenzylalcohol resin (Bachem, 0.54 mmole/g)) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

20

EXAMPLE 99

Preparation of C-terminal carboxylic acid peptides
corresponding to the above C-terminal amide sequences for
SEQ ID NOS. 62-67, 76, 77 and 81-86

25

Peptides having the sequences of SEQ ID NOS. 62-67, 76, 77 and 81-86 are assembled on the 2-chlorotritylchloride resin (200-400 mesh), 2% DVB (Novabiochem, 0.4-1.0 mmole/g)) using Fmoc-protected amino

acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 37. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides an experimentally determined (M).

EXAMPLE 100

Preparation of Peptide having SEQ ID NO. 100

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 100]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.). In general, single-coupling cycles were used throughout the synthesis and Fast Moc (HBTU activation) chemistry was employed. Deprotection (Fmoc group removal) of the growing peptide chain was achieved using piperidine. Final deprotection of the completed peptide resin was achieved using a mixture of triethylsilane (0.2 mL), ethanedithiol (0.2 mL), anisole (0.2 mL), water (0.2 mL) and trifluoroacetic acid (15 mL) according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems,

Inc.) The peptide was precipitated in ether/water (50 mL) and centrifuged. The precipitate was reconstituted in glacial acetic acid and lyophilized. The lyophilized peptide was dissolved in water). Crude purity was about 75%.

Used in purification steps and analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

The solution containing peptide was applied to a preparative C-18 column and purified (10% to 40% Solvent B in Solvent A over 40 minutes). Purity of fractions was determined isocratically using a C-18 analytical column. Pure fractions were pooled furnishing the above-identified peptide. Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 19.2 minutes. Electrospray Mass Spectrometry (M): calculated 3171.6; found 3172.

EXAMPLE 101

Preparation of Peptide having SEQ ID NO. 101

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 101]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.),

cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.9 minutes. Electrospray Mass Spectrometry (M): calculated 3179.6; found 3180.

5

EXAMPLE 102Preparation of Peptide having SEQ ID NO. 102

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 102]

15

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 37% to 47% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 12.2 minutes. Electrospray Mass Spectrometry (M): calculated 3251.6; found 3253.3.

20

25

EXAMPLE 103Preparation of Peptide having SEQ ID NO. 103

5 His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 103]

The above amidated peptide was assembled on 4-(2'-4'-
10 dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide
norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to Example 100. Used in analysis were Solvent
15 A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).
Analytical RP-HPLC (gradient 35% to 45% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide gave
product peptide having an observed retention time of 16.3
minutes. Electrospray Mass Spectrometry (M): calculated
20 3193.6; found 3197.

EXAMPLE 104Preparation of Peptide having SEQ ID NO. 104

25 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 104]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3228.6.

EXAMPLE 105

Preparation of Peptide having SEQ ID NO. 105

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 105]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3234.7.

5

EXAMPLE 106Preparation of Peptide having SEQ ID NO. 106

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂,
10 [SEQ. ID. NO. 106]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3308.7.

25

EXAMPLE 107Preparation of Peptide having SEQ ID NO. 107

His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂.

[SEQ. ID. NO. 107]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3250.7

15

EXAMPLE 108

Preparation of Peptide having SEQ ID NO. 108

His Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met
20 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 108]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

25

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
5 calculated 3252.6.

EXAMPLE 109

Preparation of Peptide having SEQ ID NO. 109

10 Ala Ala Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 109]

The above-identified amidated peptide is assembled on
15 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
20 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
25 calculated 3200.6.

EXAMPLE 110Preparation of Peptide having SEQ ID NO. 110

Ala Ala Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
5 [SEQ. ID. NO. 110]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
10 mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
15 in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3143.5.

20

EXAMPLE 111Preparation of Peptide having SEQ ID NO. 111

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 111]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
5 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
10 calculated 3214.6.

EXAMPLE 112

Preparation of Peptide having SEQ ID NO. 112

15 Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 112]

The above-identified amidated peptide is assembled on
20 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
25 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3157.5.

EXAMPLE 113

Preparation of Peptide having SEQ ID NO. 113

5

Ala Gly Asp Gly Ala Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 113]

10

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
15 purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
20 product peptide. Electrospray Mass Spectrometry (M):
calculated 3184.6.

EXAMPLE 114

Preparation of Peptide having SEQ ID NO. 114

25

Ala Gly Asp Gly Ala Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 114]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3127.5.

15

EXAMPLE 115Preparation of Peptide having SEQ ID NO. 115

Ala Gly Asp Gly Thr NaphthylAla Thr Ser Asp Leu Ser Lys
Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu
Lys Asn-NH₂ [SEQ. ID. NO. 115]

20

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

25

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3266.4.

5

EXAMPLE 116Preparation of Peptide having SEQ ID NO. 116

Ala Gly Asp Gly Thr Naphthylala Thr Ser Asp Leu Ser Lys
10 Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu
Lys Asn-NH₂ [SEQ. ID. NO. 116]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
15 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3209.4.

25

EXAMPLE 117Preparation of Peptide having SEQ ID NO. 117

Ala Gly Asp Gly Thr Phe Ser Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 117]

5 The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
10 purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
15 product peptide. Electrospray Mass Spectrometry (M):
calculated 3200.6.

EXAMPLE 118

Preparation of Peptide having SEQ ID NO. 118

20

Ala Gly Asp Gly Thr Phe Ser Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 118]

25

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and

purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3143.5.

EXAMPLE 119

Preparation of Peptide having SEQ ID NO. 119

Ala Gly Asp Gly Thr Phe Thr Ala Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 119]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3198.6.

EXAMPLE 120Preparation of Peptide having SEQ ID NO. 120

Ala Gly Asp Gly Thr Phe Thr Ala Asp Leu Ser Lys Gln Leu
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 120]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
15 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3141.5.

20

EXAMPLE 121Preparation of Peptide having SEQ ID NO. 121

Ala Gly Asp Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 121]

The above-identified peptide is assembled on 4-(2'-
4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide

norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3170.6.

EXAMPLE 122

Preparation of Peptide having SEQ ID NO. 122

Ala Gly Asp Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 122]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3113.5.

EXAMPLE 123

Preparation of Peptide having SEQ ID NO. 123

5

Ala Gly Asp Gly Thr Phe Thr Ser Glu Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 123]

10

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
15 purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
20 product peptide. Electrospray Mass Spectrometry (M):
calculated 3228.6.

EXAMPLE 124

Preparation of Peptide having SEQ ID NO. 124

25

Ala Gly Asp Gly Thr Phe Thr Ser Glu Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 124]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3171.6.

EXAMPLE 125

Preparation of Peptide having SEQ ID NO. 125

Ala Gly Asp Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 125]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3172.5.

5

EXAMPLE 126

Preparation of Peptide having SEQ ID NO. 126

Ala Gly Asp Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 126]

The above-identified amidated peptiden is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
15 acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3115.4.

25

EXAMPLE 127Preparation of Peptide having SEQ ID NO. 127

Ala Gly Asp Gly Thr Phe Thr Ser Asp Pentylgly Ser Lys Gln
5 Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys
Asn-NH₂ [SEQ. ID. NO. 127]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
15 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3230.4.

20

EXAMPLE 128Preparation of Peptide having SEQ ID NO. 128

Ala Gly Asp Gly Thr Phe Thr Ser Asp Pentylgly Ser Lys Gln
25 Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys
Asn-NH₂ [SEQ. ID. NO. 128]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3198.6.

EXAMPLE 129

Preparation of Peptide having SEQ ID NO. 129

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ala Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 129]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3141.5.

EXAMPLE 130

Preparation of Peptide having SEQ ID NO. 130

5

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ala Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 130]

10

15

20

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3157.5.

EXAMPLE 131

Preparation of Peptide having SEQ ID NO. 131

25

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Ala Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 131]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3100.4.

EXAMPLE 132

Preparation of Peptide having SEQ ID NO. 132

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Ala Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 132]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3157.6.

5

EXAMPLE 133

Preparation of Peptide having SEQ ID NO. 133

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Ala Met
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 133]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
15 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3100.5.

25

EXAMPLE 134

Preparation of Peptide having SEQ ID NO. 134

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Ala Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 134]

5 The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
10 purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
15 product peptide. Electrospray Mass Spectrometry (M):
calculated 3100.5.

EXAMPLE 135

Preparation of Peptide having SEQ ID NO. 135

20

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Ala
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 135]

25

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and

purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
5 is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3154.5.

EXAMPLE 136

10 Preparation of Peptide having SEQ ID NO. 136

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Ala
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 136]

15

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
20 Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
25 is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3115.5.

EXAMPLE 137Preparation of Peptide having SEQ ID NO. 137

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln
5 Pentylgly Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu
Lys Asn-NH₂ [SEQ. ID. NO. 137]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3212.4.

20

EXAMPLE 138Preparation of Peptide having SEQ ID NO. 138

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln
25 Pentylgly Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu
Lys Asn-NH₂ [SEQ. ID. NO. 138]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
5 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
10 calculated 3173.4.

EXAMPLE 139

Preparation of Peptide having SEQ ID NO. 139

15 Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Ala Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 139]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
20 acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
25 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
calculated 3156.6.

EXAMPLE 140Preparation of Peptide having SEQ ID NO. 140

5 Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Ala Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 140]

10 The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
15 purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
20 calculated 3099.5.

EXAMPLE 141Preparation of Peptide having SEQ ID NO. 141

25 Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Ala Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 141]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3156.6.

EXAMPLE 142

Preparation of Peptide having SEQ ID NO. 142

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Ala Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 142]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the

product peptide. Electrospray Mass Spectrometry (M):
calculated 3099.5.

EXAMPLE 143

5 Preparation of Peptide having SEQ ID NO. 143

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Ala Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 143]

10

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
15 Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
20 is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3156.6.

EXAMPLE 144

25 Preparation of Peptide having SEQ ID NO. 144

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Ala Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 144]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3099.5.

15

EXAMPLE 145Preparation of Peptide having SEQ ID NO. 145

20

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Ala Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 145]

25

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3186.6.

5

EXAMPLE 146Preparation of Peptide having SEQ ID NO. 146

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Ala Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
10 [SEQ. ID. NO. 146]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
20 in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3129.5.

25

EXAMPLE 147Preparation of Peptide having SEQ ID NO. 147

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Ala Leu Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 147]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3129.5.

15

EXAMPLE 148Preparation of Peptide having SEQ ID NO. 148

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Ala Leu Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 148]

20

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

25

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3072.4.

5

EXAMPLE 149Preparation of Peptide having SEQ ID NO. 149

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
10 Glu Glu Glu Ala Val Arg Ala Phe Ile Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 149]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
15 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3172.5.

25

EXAMPLE 150Preparation of Peptide having SEQ ID NO. 150

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
5 Glu Glu Glu Ala Val Arg Ala Phe Ile Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 150]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3115.5.

20

EXAMPLE 151Preparation of Peptide having SEQ ID NO. 151

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
25 Glu Glu Glu Ala Val Arg Leu Naphthylala Ile Glu Trp Leu
Lys Asn-NH₂ [SEQ. ID. NO. 151]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
5 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
10 calculated 3266.4.

EXAMPLE 152

Preparation of Peptide having SEQ ID NO. 152

15 Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Naphthylala Ile Glu Phe Leu
Lys Asn-NH₂ [SEQ. ID. NO. 152]

The above-identified amidated peptide is assembled on
20 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
25 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3209.4.

EXAMPLE 153

Preparation of Peptide having SEQ ID NO. 153

5

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Val Glu Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 153]

10

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
15 purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
20 product peptide. Electrospray Mass Spectrometry (M):
calculated 3200.6.

EXAMPLE 154

Preparation of Peptide having SEQ ID NO. 154

25

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Val Glu Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 154]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3143.5.

15

EXAMPLE 155Preparation of Peptide having SEQ ID NO. 155

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe tButylgly Glu Trp Leu Lys
Asn-NH₂ [SEQ. ID. NO. 155]

20

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

25

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3216.5.

5

EXAMPLE 156

Preparation of Peptide having SEQ ID NO. 156

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
10 Glu Glu Glu Ala Val Arg Leu Phe tButylgly Glu Phe Leu Lys
Asn-NH₂ [SEQ. ID. NO. 156]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
15 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3159.4.

25

EXAMPLE 157Preparation of Peptide having SEQ ID NO. 157

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Asp Trp Leu Lys Asn-NH₂
[SEQ. ID. NO. 157]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3200.6.

20

EXAMPLE 158Preparation of Peptide having SEQ ID NO. 158

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Asp Phe Leu Lys Asn-NH₂
[SEQ. ID. NO. 158]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
5 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
10 calculated 3143.5.

EXAMPLE 159

Preparation of Peptide having SEQ ID NO. 159

15 Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Ala Leu Lys Asn-NH₂
[SEQ. ID. NO. 159]

The above-identified amidated peptide is assembled on
20 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
25 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3099.5.

EXAMPLE 160

Preparation of Peptide having SEQ ID NO. 160

5

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Ala Leu Lys Asn-NH₂
[SEQ. ID. NO. 160]

10

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
15 purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
20 product peptide. Electrospray Mass Spectrometry (M):
calculated 3081.4.

EXAMPLE 161

Preparation of Peptide having SEQ ID NO. 161

25

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Ala Lys Asn-NH₂
[SEQ. ID. NO. 161]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3172.5.

15

EXAMPLE 162Preparation of Peptide having SEQ ID NO. 162

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Ala Lys Asn-NH₂
[SEQ. ID. NO. 162]

20

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

25

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3115.5.

5

EXAMPLE 163Preparation of Peptide having SEQ ID NO. 163

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Ala Asn-NH₂
10 [SEQ. ID. NO. 163]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
20 in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3157.5.

25

EXAMPLE 164Preparation of Peptide having SEQ ID NO. 164

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Ala Asn-NH₂

[SEQ. ID. NO. 164]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3100.4.

15

EXAMPLE 165

Preparation of Peptide having SEQ ID NO. 165

Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
20 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Ala-NH₂
[SEQ. ID. NO. 165]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

25

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
5 calculated 3171.6.

EXAMPLE 166

Preparation of Peptide having SEQ ID NO. 166

10 Ala Gly Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Ala-NH₂
[SEQ. ID. NO. 166]

The above-identified amidated peptide is assembled on
15 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
20 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
25 calculated 3114.5.

EXAMPLE 167Preparation of Peptide having SEQ ID NO. 167

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH₂ [SEQ. ID. NO.
167]

The above-identified amidated peptide is assembled on
10 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
20 calculated 4033.5.

EXAMPLE 168Preparation of Peptide having SEQ ID NO. 168

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro-NH₂ [SEQ. ID. NO.
168]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3984.4.

EXAMPLE 169

Preparation of Peptide having SEQ ID NO. 169

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro-NH₂ [SEQ. ID. NO. 169]

20

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide

25

is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 4016.5.

5

EXAMPLE 170Preparation of Peptide having SEQ ID NO. 170

His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
10 Gly Gly Pro Ser Ser Gly Ala Pro-NH₂ [SEQ. ID. NO. 170]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
20 in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3861.3.

25

EXAMPLE 171Preparation of Peptide having SEQ ID NO. 171

Ala Gly Glu Gly Thr Phe Thr Ser Asp Ala Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn

Gly Gly Pro Ser Ser Gly Ala Pro-NH₂ [SEQ. ID. NO. 171]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3746.1.

15

EXAMPLE 172

Preparation of Peptide having SEQ ID NO. 172

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
20 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 172]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA

25

in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
5 calculated 3742.1.

EXAMPLE 173

Preparation of Peptide having SEQ ID NO. 173

10 His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 173]

The above-identified amidated peptide is assembled on
15 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
20 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
25 calculated 3693.1.

EXAMPLE 174Preparation of Peptide having SEQ ID NO. 174

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly-NH₂ [SEQ. ID. NO. 174]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3751.2.

20

EXAMPLE 175Preparation of Peptide having SEQ ID NO. 175

His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser-NH₂ [SEQ. ID. NO. 175]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
5 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
10 calculated 3634.1.

EXAMPLE 176

Preparation of Peptide having SEQ ID NO. 176

15 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser-NH₂ [SEQ. ID. NO. 176]

The above-identified amidated peptide is assembled on
20 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
25 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):

calculated 3526.9.

EXAMPLE 177

Preparation of Peptide having SEQ ID NO. 177

5

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser-NH₂ [SEQ. ID. NO. 177]

10

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
15 purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
20 product peptide. Electrospray Mass Spectrometry (M):
calculated 3477.9.

EXAMPLE 178

Preparation of Peptide having SEQ ID NO. 178

25

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro-NH₂ [SEQ. ID. NO. 178]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3519.9.

15

EXAMPLE 179Preparation of Peptide having SEQ ID NO. 179

His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly-NH₂ [SEQ. ID. NO. 179]

20

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B

25

in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3307.7.

5

EXAMPLE 180

Preparation of Peptide having SEQ ID NO. 180

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
10 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly-NH₂ [SEQ. ID. NO. 180]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
15 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
20 in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3186.5.

25

EXAMPLE 181

Preparation of Peptide having SEQ ID NO. 181

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly tPro Ser Ser Gly Ala tPro tPro tPro-NH₂ [SEQ. ID.
NO. 181]

5

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
10 Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Double couplings
are required at residues 37,36 and 31. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
15 in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 4121.1.

20

EXAMPLE 182

Preparation of Peptide having SEQ ID NO. 182

His Gly Glu Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala tPro tPro tPro-NH₂ [SEQ. ID.
25 NO. 182].

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55

mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Double couplings are required at residues 37, 36 and 31. Used in analysis
5 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
10 calculated 4173.2.

EXAMPLE 183

Preparation of Peptide having SEQ ID NO. 183

15 His Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly NMeala Ser Ser Gly Ala NMeala NMeala-NH₂, [SEQ. ID.
NO. 183]

20 The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
25 purified in a similar way to Compound 1. Double couplings
are required at residues 36 and 31. Used in analysis are
Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in
ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in
Solvent A over 30 minutes) of the lyophilized peptide is

then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3796.1.

5

EXAMPLE 184Preparation of Peptide having SEQ ID NO. 184

Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
10 Gly Gly hPro Ser Ser Gly Ala hPro-NH₂ [SEQ. ID. NO. 184]

The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. A double coupling is required at residue 31. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in
20 ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M): calculated 3871.1.

25

EXAMPLE 185Preparation of Peptide having SEQ ID NO. 185

His Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
5 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala-NH₂ [SEQ. ID. NO. 185]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
10 acetamide norleucine MBHA resin (Novabiochem, 0.55
mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
15 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 3750.2.

20

EXAMPLE 186Preparation of Peptide having SEQ ID NO. 186

His Gly Asp Ala Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
25 Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly-NH₂ [SEQ. ID. NO. 186]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy

acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Example 100. Used in analysis
5 are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry (M):
10 calculated 3408.8.

EXAMPLE 187

Preparation of Peptide having SEQ ID NO. 187

15 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH₂ [SEQ. ID.
NO. 187]

20 The above-identified amidated peptide is assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and
25 purified in a similar way to Example 100. Used in analysis are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the

product peptide. Electrospray Mass Spectrometry (M):
calculated 4120.6.

EXAMPLE 188

5 Preparation of Peptide having SEQ ID NO. 188

Ala Gly Ala Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu
Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn
Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH₂ [SEQ. ID.
10 NO. 188]

The above-identified amidated peptide is assembled on
4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy
acetamide norleucine MBHA resin (Novabiochem, 0.55
15 mmole/g) using Fmoc-protected amino acids (Applied
Biosystems, Inc.), cleaved from the resin, deprotected and
purified in a similar way to Example 100. Used in analysis
are Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA
in ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B
20 in Solvent A over 30 minutes) of the lyophilized peptide
is then carried out to determine the retention time of the
product peptide. Electrospray Mass Spectrometry (M):
calculated 4005.5.

25 EXAMPLE 189

Preparation of C-terminal carboxylic acid peptides
corresponding to the above C-terminal amide sequences for
Peptides having SEQ ID NOS. 100-166, 172-177, 179-180 and
185-188.

C-terminal carboxylic acid peptides corresponding to
amidated having SEQ ID NOS. 100-166, 172-177, 179-180 and
185-188 are assembled on the so called Wang resin (p-
5 alkoxybenzylalcohol resin (Bachem, 0.54 mmole/g)) using
Fmoc-protected amino acids (Applied Biosystems, Inc.),
cleaved from the resin, deprotected and purified in a
similar way to that described in Example 100. Used in
analysis are Solvent A (0.1% TFA in water) and Solvent B
10 (0.1% TFA in ACN). Analytical RP-HPLC (gradient 30% to 60%
Solvent B in Solvent A over 30 minutes) of the lyophilized
peptide is then carried out to determine the retention
time of the product peptide. Electrospray Mass
Spectrometry provides an experimentally determined (M).

15

EXAMPLE 190

Preparation of C-terminal carboxylic acid peptides
corresponding to the above C-terminal amide sequences for
20 Peptides having SEQ ID NOS. 167-171, 178 and 181-184.

C-terminal carboxylic acid peptides corresponding to
amidated SEQ ID NOS. 167-171, 178 and 181-184 are
assembled on the 2-chlorotritylchloride resin (200-400
25 mesh), 2% DVB (Novabiochem, 0.4-1.0 mmole/g)) using Fmoc-
protected amino acids (Applied Biosystems, Inc.), cleaved
from the resin, deprotected and purified in a similar way
to that described in Example 100. Used in analysis are
Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in

ACN). Analytical RP-HPLC (gradient 30% to 60% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide is then carried out to determine the retention time of the product peptide. Electrospray Mass Spectrometry provides
5 an experimentally determined (M).

Various modifications of the invention in addition to those shown and described herein will become apparent to
10 those skilled in the art from the foregoing description and fall within the scope of the following claims.

WE CLAIM:

1. A method for treating conditions or disorders which can be alleviated by reducing food intake in a subject comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist.
2. The method according to claim 1 wherein said exendin or exendin agonist is administered parenterally.
3. The method according to claim 2 wherein said parenteral administration is by injection.
4. The method according to claim 3 wherein the injection is a peripheral injection.
5. The method according to claim 1 wherein about 10 μ g-30 μ g to about 5mg of the exendin or exendin agonist is administered per day.
6. The method according to claim 1 wherein about 10 μ g-30 μ g to about 2 mg of the exendin or exendin agonist is administered per day.
7. The method according to claim 1, wherein about 30 μ g to about 500 μ g of the exendin or exendin agonist is administered per day.
8. The method of claim 1 wherein said condition or disorder is obesity.
9. The method of claim 1 wherein said condition or disorder is Type II diabetes.
10. The method of claim 1 wherein said subject is human.
11. The method of claim 1 wherein said condition or disorder is an eating disorder.

12. The method of claim 1 wherein said condition or disorder is insulin-resistance syndrome.

13. A method for reducing the appetite of a subject comprising administering to said subject an appetite-
5 lowering amount of an exendin or an exendin agonist.

14. A method for reducing the weight of a subject comprising administering to said subject a therapeutically effective amount of an exendin or an exendin agonist.

15. A method for lowering plasma lipids comprising
10 administering to said subject a therapeutically effective amount of an exendin or an exendin agonist.

16. The method according to any of claims 1-15 wherein said exendin is exendin-3.

17. The method according to any of claims 1-15
15 wherein said exendin is exendin-4.

18. The method according to any of claims 1-15 wherein said exendin agonist is selected from the group consisting of exendin-4 (1-30), exendin-4 (1-30) amide, exendin-4 (1-28) amide, ¹⁴Leu, ²⁵Phe exendin-4 amide,
20 ¹⁴Leu, ²⁵Phe exendin-4 (1-28) amide, and ¹⁴Leu, ²²Ala, ²⁵Phe exendin-4 (1-28) amide.

19. The method according to any of claims 1-15, further comprising administering a therapeutically effective amount of one or more compounds selected from the
25 group consisting essential of an amylin agonist, a leptin, and a CCK.

20. The method according to any of claims 1-15 wherein said exendin agonist is an exendin agonist according to Formula I.

21. The method according to any of claims 1-15 wherein said exendin agonist is an exendin agonist according to Formula II.

5 22. The method according to any of claims 1-15 wherein said exendin agonist is an exendin agonist according to Formula III.

10 23. A pharmaceutical composition for use in the treatment of conditions or disorders associated with hypernutrition comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

24. The pharmaceutical composition according to claim 21, wherein said exendin is exendin-3.

15 25. The pharmaceutical composition according to claim 21 wherein said exendin is exendin-4.

26. The pharmaceutical composition according to claim 21 wherein said exendin agonist is selected from the group consisting of exendin-4 (1-30), exendin-4 (1-30) amide, exendin-4 (1-28) amide, ¹⁴Leu, ²⁵Phe exendin-4 amide, 20 ¹⁴Leu, ²⁵Phe exendin-4 (1-28) amide, and ¹⁴Leu, ²²Ala, ²⁵Phe exendin-4 (1-28) amide.

27. The pharmaceutical composition of claim 21 wherein said therapeutically effective amount is a therapeutically effective amount for a human subject.

25 28. A pharmaceutical composition for use in reducing the appetite of a subject comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.


29. A pharmaceutical composition for use in reducing the weight of a subject comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

5 30. A pharmaceutical composition for use in lowering the plasma lipid level of a subject comprising a therapeutically effective amount of an exendin or exendin agonist in association with a pharmaceutically acceptable carrier.

10 31. The pharmaceutical composition according to any of claims 21-28, further comprising a therapeutically effective amount of one or more compounds selected from the group consisting essentially of an amylin agonist, a leptin, and a CCK.


1/11

FIG. 1.






2/11

FIG. 2.




3/11

FIG. 3.




4/11

FIG. 4.




5/11

FIG. 5.




6/11

FIG. 6.




7/11

FIG. 7.




8/11

FIG. 8.



9/11

FIG. 9.



10/11

1 15 20
 Xaa₁ Xaa₂ Xaa₃ Gly Thr Xaa₄ Xaa₅ Xaa₆ Xaa₇ Xaa₈ Ser Lys Gln Xaa₉ Glu Glu Glu Ala Val Arg Leu
 25 30 35
 Xaa₁₀ Xaa₁₁ Xaa₁₂ Xaa₁₃ Leu Lys Asn Gly Gly Xaa₁₄ Ser Ser Gly Ala Xaa₁₅ Xaa₁₆ Xaa₁₇ Xaa₁₈-Z

[SEQ. ID. NO.]	Xaa ₁	Xaa ₂	Xaa ₃	Xaa ₄	Xaa ₅	Xaa ₆	Xaa ₇	Xaa ₈	Xaa ₉	Xaa ₁₀	Xaa ₁₁	Xaa ₁₂	Xaa ₁₃	Xaa ₁₄	Xaa ₁₅	Xaa ₁₆	Xaa ₁₇	Xaa ₁₈	Z
9	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH ₂
10	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
11	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH ₂
12	Tyr	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
13	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Tyr	NH ₂
14	His	Gly	Asp	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
15	His	Gly	Glu	naph	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
16	His	Gly	Glu	Phe	Ser	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
17	His	Gly	Glu	Phe	Ser	Thr	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
18	His	Gly	Glu	Phe	Thr	Thr	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
19	His	Gly	Glu	Phe	Thr	Ser	Glu	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
20	His	Gly	Glu	Phe	Thr	Ser	Asp	pGly	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
21	His	Gly	Glu	Phe	Thr	Ser	Asp	pGly	Leu	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH ₂
22	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	pGly	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂

FIGURE 10
 (Sheet 1 of 2)

11/11

[SEQ. ID. NO.]	Xaa ₁	Xaa ₂	Xaa ₃	Xaa ₄	Xaa ₅	Xaa ₆	Xaa ₇	Xaa ₈	Xaa ₉	Xaa ₁₀	Xaa ₁₁	Xaa ₁₂	Xaa ₁₃	Xaa ₁₄	Xaa ₁₅	Xaa ₁₆	Xaa ₁₇	Xaa ₁₈	Z
23	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	pGly	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH ₂
24	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	naph	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
25	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Val	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
26	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Val	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH
27	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	tBuG	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
28	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	tBuG	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH ₂
29	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Asp	Trp	Pro	Pro	Pro	Pro	Ser	NH ₂
30	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH ₂
31	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	tPro	tPro	tPro	tPro	Ser	NH ₂
32	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	tPro	tPro	tPro	Ser	NH ₂
33	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	hPro	hPro	hPro	hPro	Ser	NH ₂
34	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	hPro	hPro	hPro	Ser	NH ₂
35	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	tPro	tPro	tPro	tPro	Ser	NH ₂
36	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	hPro	hPro	hPro	hPro	Ser	NH ₂
37	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	MeAla	MeAla	MeAla	MeAla	Ser	NH
38	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	MeAla	MeAla	MeAla	Ser	NH
39	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	MeAla	MeAla	MeAla	MeAla	Ser	NH

FIGURE 10
(Sheet 2 of 2)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/00449

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 38/16

US CL : 514/2, 866

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/2, 866

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS ONLINE, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,424,286 A (ENG) 13 June 1995 see entire document.	1-31

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

07 MAY 1998

Date of mailing of the international search report

29 MAY 1998

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Zohreh Fay

Telephone No. (703) 308-1235

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.